

bsm-51558M**[Primary Antibody]****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

UCHL1/PGP9.5 Mouse mAb**— DATASHEET —**


Host: Mouse	Isotype: IgG1	Applications: WB (1:500-1000) IHC-P (1:20-50) IHC-F (1:400-800) IF (1:100-500) Reactivity: Human (predicted: Mouse, Rat, Pig, Cow, Monkey, Xenopus) Predicted MW.: 25 kDa Subcellular Location: Cell membrane ,Cytoplasm
Clonality: Monoclonal	CloneNo.: T2F5	
GeneID: 7345	SWISS: P09936	
Target: UCHL1/PGP9.5		
Immunogen: KLH conjugated synthetic peptide derived from human UCHL1/PGP9.5: 151-223/223.		
Purification: affinity purified by Protein G		
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: catalyzes the hydrolysis of ubiquitin carboxy-terminal thiolesters to form ubiquitin and a thiol; may play a role in neuropathic pain [RGD]. Found in neuronal cell bodies and processes throughout the neocortex (at protein level). Expressed in neurons and cells of the diffuse neuroendocrine system and their tumors. Weakly expressed in ovary. Down-regulated in brains from Parkinson disease and Alzheimer disease patients.		

— VALIDATION IMAGES —

Sample: Lane 1: 293T cell lysates Lane 2: NCI-H1299 cell lysates Lane 3: PC-12 cell lysates
Primary: Anti-UCHL1/PGP9.5 (bsm-51558M) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Mouse IgG at 1/20000 dilution Predicted band size: 25 kD Observed band size: 25 kD



Paraformaldehyde-fixed, paraffin embedded (human astrogloma section); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with UCHL1/PGP9.5) Monoclonal Antibody, Unconjugated (bsm-51558M) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Mouse)(sp-0024) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (human brain section); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with UCHL1/PGP9.5) Monoclonal Antibody, Unconjugated (bsm-51558M) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Mouse)(sp-0024) instructions and DAB staining.

— SELECTED CITATIONS —

- **[IF=1.1]** Alkotami Ahmed S.. et al. Diabetic small fiber neuropathy: clinical and electrophysiological study. EGYPT J NEUROL PSYCH. 2024 Dec;60(1):1-16 IHC ;Human. 10.1186/s41983-024-00923-8