bsm-51591M

[Primary Antibody]

Bioss ANTIBODIES

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

phospho-RPS6 (Ser235/Ser236) Mouse mAb

- DATASHEET -

Host: Mouse Isotype: IgG1
Clonality: Monoclonal CloneNo.: J88A1
GeneID: 6194 SWISS: P62753

Target: RPS6 (Ser235/Ser236) **Purification:** affinity purified by Protein G

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: Ribosomes, the organelles that catalyze protein synthesis, consist

of a small 40S subunit and a large 60S subunit. Together these subunits are composed of 4 RNA species and approximately 80 structurally distinct proteins. This gene encodes a cytoplasmic ribosomal protein that is a component of the 40S subunit. The protein belongs to the S6E family of ribosomal proteins. It is the major substrate of protein kinases in the ribosome, with subsets of five C-terminal serine residues phosphorylated by different protein kinases. Phosphorylation is induced by a wide range of stimuli, including growth factors, tumor-promoting agents, and mitogens. Dephosphorylation occurs at growth arrest. The protein may contribute to the control of cell growth and proliferation through the selective translation of particular classes of mRNA. As is typical for genes encoding ribosomal proteins, there are multiple processed pseudogenes of this gene dispersed through the

genome. [provided by RefSeq, Jul 2008]

Applications: WB (1:500-2000)


Reactivity: Human

Predicted MW.: ^{29 kDa}

Subcellular Cell membrane, Cytoplasm

Location: , Nucleus

- VALIDATION IMAGES -



MCF-7 (H) Serum starvation overnight, then grown in 20% FBS media for 30 min, 25 µg total protein per lane of cell lysates (see on figure) probed with Phospho-RPS6 (Ser235/Ser236) monoclonal antibody, unconjugated (bsm-51591M) at 1:1000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min.