bsm-51659M

[Primary Antibody]

BIOSS ANTIBODIES

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

Hexokinase II Mouse mAb

- DATASHEET -

Host: Mouse Isotype: IgG1, k
Clonality: Monoclonal CloneNo.: S2G6
GeneID: 3099 SWISS: P52789

Target: Hexokinase II

Purification: affinity purified by Protein G

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: The hexokinases utilize Mg-ATP as a phosphoryl donor to catalyze the first step of intracellular glucose metabolism, the conversion of

glucose to glucose-6-phosphate. Four hexokinase isoenzymes have been identified, including hexokinase I (HXK I), hexokinase II (HXK II), hexokinase II (HXK III), hexokinase III (HXK III) and hexokinase IV (HXK IV, also designated glucokinase or GCK). Hexokinases I-III each contain an N-terminal cluster of hydrophobic amino acids. Glucokinase lacks the N-terminal hydrophobic cluster. The hydrophobic cluster is thought to be necessary for membrane binding. This is substantiated by the finding that glucokinase has lower affinity for glucose than do the other hexokinases. HXK I has been shown to be

glucose than do the other hexokinases. HXK I has been shown to be expressed in brain, kidney and heart tissues as well as in hepatoma cell lines. HXK II is involved in the uptake and utilization of glucose by adipose and skeletal tissues. Of the hexokinases, HXK III has the highest affinity for glucose. Glucokinase is expressed in pancreatic beta cells where it functions as a glucose sensor, determining the

"set point" for insulin secretion.

Applications: WB (1:500-1000)

IHC-P (1:50-200) IHC-F (1:50-200) IF (1:50-200)

Reactivity: Human

Predicted MW.: 101 kDa

Subcellular Location: Cell membrane ,Cytoplasm

- VALIDATION IMAGES -



Sample: Lane 1: Hela cell lysates Lane 2: 293 cell lysates Lane 3: Jurkat cell lysates Primary: Anti-Hexokinase II (bsm-51659M) at 1/500~1000 dilution Secondary: IRDye800CW Goat Anti-Mouse IgG at 1/20000 dilution Predicted band size: 101 kD Observed band size: 101 kD