bsm-52478R

[Primary Antibody]

Bioss ANTIBODIES

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

Collagen I Recombinant Rabbit mAb

- DATASHEET -

Host: Rabbit Isotype: IgG
Clonality: Recombinant CloneNo.: 5F2
GeneID: 1277 SWISS: P02452

Target: Collagen I

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: Collagens are highly conserved throughout evolution and are

characterised by an uninterrupted "Glycine X Y" triplet repeat that is a necessary part of the triple helical structure. Type I collagen (95 kDa) is found in bone, cornea, skin and tendon. Mutations in the encoding gene are associated with osteogenesis imperfecta, Ehlers Danlos syndrome, and idiopathic osteoporosis. Reciprocal translocations between chromosomes 17 and 22, where this gene and the gene for Platelet-derived growth factor beta are located, are associated with a particular type of skin tumor called dermatofibrosarcoma protuberans, resulting from unregulated

expression of the growth factor.

Applications: IHC-P (1:100-500)

IHC-F (1:100-500) IF (1:100-500) Flow-Cyt (1:50-100)


Reactivity: Human (predicted: Cow)

Predicted MW.: 130 kDa


Subcellular Secreted ,Extracellular

Location: matrix


VALIDATION IMAGES



Paraformaldehyde-fixed, paraffin embedded Human Breast Cancer; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with Collagen I Monoclonal Antibody, Unconjugated (bsm-52478R) at 1:200 overnight at 4°C, followed by conjugation to the bs-0295G-HRP and DAB (C-0010) staining.



Paraformaldehyde-fixed, paraffin embedded Human Uterus; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with Collagen I Monoclonal Antibody, Unconjugated (bsm-52478R) at 1:200 overnight at 4°C, followed by conjugation to the bs-0295G-HRP and DAB (C-0010) staining.



The U-87MG (H) cells were fixed with 4% PFA (10 min at r.t.) and then permeabilized with 90% ice-cold methanol for 20 min at -20°C, the cells then were incubated in 5%BSA to block non-specific protein-protein interactions (30 min at r.t.). Primary Antibody (green): Rabbit Anti-Collagen I antibody (bsm-52478R,1:100); Secondary Antibody (white blue): Goat anti-Rabbit IgG-BF488(bs-60295G-BF488): 1 μ g/test. Blank control (black): PBS. Acquisition of 20,000 events was performed.

- SELECTED CITATIONS -

- [IF=11.322] Jinxiu Yu. et al. Promoting osseointegration of titanium by pH-responsive releasing of H2S and optimizing polarization time for macrophages. COMPOS PART B-ENG. 2023 Mar;253:110554 IHC; Rat. 10.1016/j.compositesb.2023.110554
- [IF=3.913] Xiaoliang Zhou. et al. Ursolic acid inhibits human dermal fibroblasts hyperproliferation, migration, and collagen deposition induced by TGF-β via regulating the Smad2/3 pathway. GENE. 2023 May;867:147367 WB; Human. 36931410
- [IF=3.2] Yiyang Fan. et al. Effects and Significance of Dicliptera chinensis Polysaccharide on the Expression of

Transforming Growth Factor β 1/Connective Tissue Growth Factor Pathway in the Masseter and Head and Neck Skin of Rats With Radiation-Induced Fibrosis. INT DENT J. 2024 Jul;: IHC; Rat. 38991877

- [IF=3.2] Yiyang Fan. et al.Effects and Significance of Dicliptera chinensis Polysaccharide on the Expression of Transforming Growth Factor β1/Connective Tissue Growth Factor Pathway in the Masseter and Head and Neck Skin of Rats With Radiation-Induced Fibrosis..INTERNATIONAL DENTAL JOURNAL.2025 Apr;75(2):784-796. IHC; Rat. 38991877
- [IF=2.6] Han Wang. et al. Ganoderma lucidum extract reduces skin aging by reducing mitochondrial stress and controlling mitochondrial numbers. FITOTERAPIA. 2025 May;:106627 IHC; Mouse. 40381851