
MX1 Mouse mAb

Catalog Number: bsm-51528M

Target Protein: MX1

Concentration: 1mg/ml

Form: Liquid

Host: Mouse

Clonality: Monoclonal

Clone No.: C3G6

Isotype: IgG1

Applications: WB (1:500-1000)

Reactivity: Human (predicted:Pig)

Predicted MW: 73 kDa

Entrez Gene: 4599

Swiss Prot: P20591

Source: KLH conjugated synthetic peptide derived from human MX1: 601-662/662.

Purification: affinity purified by Protein G


Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: bs-6653P is one synthetic peptide derived from human MX1.

The Dynamin family of microtubule-associated proteins function as GTPases that are involved in microtubule bundling and endocytosis. In mice, Mx2 (myxovirus resistance protein 2) and Mx1 (myxovirus resistance protein 1) are members of the Dynamin family that are involved in the immune response to viral infections. Localized to the cytoplasm, Mx2 contains one GED domain and is expressed in response to viral infection or treatment by IFN-alpha/IFN-beta. Once expression is induced, Mx2 accumulates in the cytoplasm and inhibits the replication of vesicular stomatitis virus (VSV), thereby conferring resistance to VSV infection. Unlike Mx2, Mx1 is localized to the nucleus where, upon induction by IFN-alpha/IFN-beta, it provides selective resistance to infection by the highly lethal H5N1 influenza virus. In humans, MxA and MxB function in a similar manner to Mx1 and Mx2, conferring resistance to specific target viruses. Mx3 is a rat-specific member of the myxovirus resistance protein family.

VALIDATION IMAGES



Sample: Lane 1: T47D cell lysates Primary: Anti-MX1 (bsm-51528M) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Mouse IgG at 1/20000 dilution Predicted band size: 73 kD Observed band size: 73 kD