

**bsm-60645R****[ Primary Antibody ]****BioSS**  
**ANTIBODIES**

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

**phospho-AKT (Thr308) Recombinant Rabbit mAb****— DATASHEET —**

<b>Host:</b> Rabbit	<b>Isotype:</b> IgG	<b>Applications:</b> WB (1:500-2000)
<b>Clonality:</b> Recombinant	<b>CloneNo.:</b> R3F4	<b>Reactivity:</b> Mouse
<b>GeneID:</b> 208	<b>SWISS:</b> P31751	
<b>Target:</b> AKT (Thr308)		
<b>Purification:</b> affinity purified by Protein A		<b>Predicted MW.:</b> 56 kDa
<b>Concentration:</b> 1mg/ml		<b>Subcellular Location:</b> Cell membrane ,Cytoplasm ,Nucleus
<b>Storage:</b> 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
<b>Background:</b> This gene is a putative oncogene encoding a protein belonging to a subfamily of serine/threonine kinases containing SH2-like (Src homology 2-like) domains. The gene was shown to be amplified and overexpressed in 2 of 8 ovarian carcinoma cell lines and 2 of 15 primary ovarian tumors. Overexpression contributes to the malignant phenotype of a subset of human ductal pancreatic cancers. The encoded protein is a general protein kinase capable of phosphorylating several known proteins. [provided by RefSeq, Jul 2008]		

**— VALIDATION IMAGES —**

Blocking buffer: 5% NFDM/TBST Primary Ab  
dilution: 1:1000 Primary Ab incubation  
condition: 2 hours at room temperature  
Secondary Ab: Goat Anti-Rabbit IgG H&L (HRP)  
Lysate: 1: NIH/3T3, 2: NIH/3T3 + PDGF (100  
ng/mL, 15 min) Protein loading quantity: 20 ug  
Exposure time: 60 s Predicted MW: 56 kDa  
Observed MW: 60 kDa

**— SELECTED CITATIONS —**

- **[IF=6.8]** Bingjie Ge. et al. Integrated network toxicology, molecular docking, and in vivo experiments to elucidate molecular mechanism of aflatoxin B1 hepatotoxicity. ECOTOX ENVIRON SAFE. 2024 Apr;275:116278 WB ;Mouse. 38564860
- **[IF=4.8]** Wang Yunyun. et al. Iridoids rich fraction from Valeriana jatamansi Jones promotes axonal regeneration and motor functional recovery after spinal cord injury through activation of the PI3K/Akt signaling pathway. FRONT MOL NEUROSCI. 2024 May;17: WB ;Rat. 38756705