

bsm-60666R**[Primary Antibody]**

Histone H3 (Di-Methyl Lys122) Recombinant Rabbit mAb

BioSS
ANTIBODIES

www.bioss.com.cn

sales@bioss.com.cn


techsupport@bioss.com.cn

400-901-9800


— DATASHEET —

Host: Rabbit	Isotype: IgG	Applications: WB (1:500-2000) Reactivity: Human, Mouse (predicted: Rat) Predicted MW.: 15 kDa Subcellular Location: Nucleus
Clonality: Recombinant	CloneNo.: R11A1	
GeneID: 8350	SWISS: P68431	
Target: Histone H3 (Di-Methyl Lys122)		
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: Modulation of the chromatin structure plays an important role in the regulation of transcription in eukaryotes. The nucleosome, made up of four core histone proteins (H2A, H2B, H3 and H4), is the primary building block of chromatin. The N-terminal tail of core histones undergoes different posttranslational modifications including acetylation, phosphorylation and methylation. These modifications occur in response to cell signal stimuli and have a direct effect on gene expression. In most species, the histone H2B is primarily acetylated at lysines 5, 12, 15 and 20. Histone H3 is primarily acetylated at lysines 9, 14, 18 and 23. Acetylation at lysine 9 appears to have a dominant role in histone deposition and chromatin assembly in some organisms. Phosphorylation at Ser10 of histone H3 is tightly correlated with chromosome condensation during both mitosis and meiosis.		

— VALIDATION IMAGES —



Blocking buffer: 5% NFDM/TBST Primary Ab
 dilution: 1:1000 Primary Ab incubation
 condition: 2 hours at room temperature
 Secondary Ab: 1: Mouse brain, 2: Rat brain, 3:
 Recombinant histone H3 Protein loading
 quantity: 20 µg Exposure time: 60 s Predicted
 MW: 15 KDa Observed MW: 15 KDa



Blocking buffer: 5% NFDM/TBST Primary Ab
 dilution: 1:1000 Primary Ab incubation
 condition: 2 hours at room temperature
 Secondary Ab: 1: HeLa cytosol, 2: HeLa nuclear
 Protein loading quantity: 20 µg Exposure time:
 60 s Predicted MW: 15 KDa Observed MW: 15 KDa