

bsm-34042M**[Primary Antibody]****Lamin A/C Mouse mAb****Bioss**
ANTIBODIES

www.bioss.com.cn

sales@bioss.com.cn


techsupport@bioss.com.cn

400-901-9800


— DATASHEET —**Host:** Mouse**Clonality:** Monoclonal**GeneID:** 4000**Target:** Lamin A/C**Immunogen:** KLH conjugated synthetic peptide derived from human lamin A: 1-100/664.**Concentration:** 1mg/ml**Storage:** 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.


Background: The nuclear lamina consists of a two-dimensional matrix of proteins located next to the inner nuclear membrane. The lamin family of proteins make up the matrix and are highly conserved in evolution. During mitosis, the lamina matrix is reversibly disassembled as the lamin proteins are phosphorylated. Lamin proteins are thought to be involved in nuclear stability, chromatin structure and gene expression. Vertebrate lamins consist of two types, A and B. Alternative splicing results in multiple transcript variants. Mutations in this gene lead to several diseases: Emery-Dreifuss muscular dystrophy, familial partial lipodystrophy, limb girdle muscular dystrophy, dilated cardiomyopathy, Charcot-Marie-Tooth disease, and Hutchinson-Gilford progeria syndrome. [provided by RefSeq, Apr 2012]

Isotype: IgG**CloneNo.:** 3E1**SWISS:** P02545**Applications:** **WB** (1:500-1000)**IHC-P** (1:50-500)**IHC-F** (1:50-500)**IF** (1:50-500)**ICC/IF** (1:50-100)**ELISA** (1:5000-10000)**Reactivity:** Human, Mouse, Rat
(predicted: Pig, Cow, Dog, Horse)**Predicted MW.:** 69/62 kDa**Subcellular Location:** Nucleus**— VALIDATION IMAGES —**


25 ug total protein per lane of various lysates (see on figure) probed with Lamin A/C monoclonal antibody, unconjugated (bsm-34042M) at 1:200 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min.



Paraformaldehyde-fixed, paraffin embedded Human Breast Cancer; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with Lamin A/C Monoclonal Antibody, Unconjugated (bsm-34042M) at 1:50 overnight at 4°C, followed by conjugation to the bs-40296G-HRP and DAB (C-0010) staining.



Paraformaldehyde-fixed, paraffin embedded Human Colon Cancer; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with Lamin A/C Monoclonal Antibody, Unconjugated (bsm-34042M) at 1:50 overnight at 4°C, followed by conjugation to the bs-40296G-HRP and DAB (C-0010) staining.



HeLa cell; 4% Paraformaldehyde-fixed; Triton X-100 at room temperature for 20 min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Antibody incubation with (Lamin A/C) monoclonal Antibody, Unconjugated (bsm-34042M) 1:100, 90 minutes at 37°C;

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

followed by a conjugated Goat Anti-Rabbit IgG antibody at 37°C for 90 minutes, DAPI (blue, C02-04002) was used to stain the cell nuclei.