bs-1754R

[Primary Antibody]

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

EAR1 Rabbit pAb

DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal GenelD: 13586 Target: EAR1

Immunogen: KLH conjugated synthetic peptide derived from mouse Eosinophil

cationic protein 1: 65-155/155.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: Eosinophil derived neurotoxin (EDN) is a protein belonging to the

ribonuclease (RNase) A superfamily. It has recently been found to have antiviral activity against respiratory syncytial virus and

human immunodeficiency virus in vitro.

Applications: WB (1:500-2000)


IHC-P (1:100-500) **IHC-F** (1:100-500) **IF** (1:100-500) Flow-Cyt (1ug/Test)

Reactivity: Human, Mouse, Rat

Predicted MW.:

Subcellular Cytoplasm ,Nucleus


VALIDATION IMAGES



Sample: Lane 1: Mouse Spleen tissue lysates Lane 2: Mouse Bone tissue lysates Primary: Anti-EAR1 (bs-1754R) at 1/200 dilution Secondary: IRDve800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 14 kDa Observed band size: 17 kDa



Paraformaldehyde-fixed, paraffin embedded (rat pancreas tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (EAR1) Polyclonal Antibody. Unconjugated (bs-1754R) at 1:400 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.



Blank control (Black line): Molt4 (Black). Primary Antibody (green line): Rabbit Anti-EAR1 antibody (bs-1754R) Dilution: 1µg/10^6 cells; Isotype Control Antibody (orange line): Rabbit IgG. Secondary Antibody (white blue line): Goat antirabbit IgG-AF647 Dilution: 1µg /test. Protocol The cells were fixed with 4% PFA (10min at room temperature) and then permeabilized with 90% ice-cold methanol for 20 min at room temperature. The cells were then incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at room temperature .Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.

— SELECTED CITATIONS —

- [IF=25.841] Xuefeng Fei. et al. Neddylation of Corola determines the fate of multivesicular bodies and biogenesis of extracellular vesicles. J Extracell Vesicles. 2021 Oct;10(12):e12153 WB; Mouse. 34623756
- [IF=5.52] Park, Shin Yong, et al. "Peptidoglycan Recognition Protein 1 Enhances Experimental Asthma by Promoting Th2 and Th17 and Limiting Regulatory T Cell and Plasmacytoid Dendritic Cell Responses." The Journal of Immunology (2013). Other; Mouse. 23420883
- [IF=2.629] Yu Wen-Yan. et al. Acupuncture Alleviates Menstrual Pain in Rat Model via Suppressing Eotaxin/CCR3 Axis to

dolescents. J ASTHM	A ALLERGY. 2023 Oct	29 IHC ;Huma	n. 37927775	