

bs-9189R**[Primary Antibody]****ZNF354C Rabbit pAb****BioSS**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

— DATASHEET —

Host: Rabbit	Isotype: IgG	Applications: IHC-P (1:100-500) IHC-F (1:100-500) IF (1:50-200) Reactivity: Rat (predicted: Human, Mouse, Cow, Dog, Horse) Predicted MW.: 64 kDa Subcellular Location: Nucleus
Clonality: Polyclonal		
GeneID: 147011	SWISS: Q8NCQ7	
Target: ZNF354C		
Immunogen: KLH conjugated synthetic peptide derived from human ZNF354C: 401-500/554.		
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: May function as a transcription repressor. Suppresses osteogenic effects of RUNX2. Binds to 5'-CCACA-3' core sequence. May be involved in osteoblastic differentiation.		

— VALIDATION IMAGES —

Paraformaldehyde-fixed, paraffin embedded (rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (ZNF354C) Polyclonal Antibody, Unconjugated (bs-9189R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (rat testis); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (ZNF354C) Polyclonal Antibody, Unconjugated (bs-9189R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.