bs-2029R

[Primary Antibody]

www.bioss.com.cn sales@bioss.com.cn

Morphine Rabbit pAb

DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Target: Morphine

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: Morphine is thought to produce reinforcement phenomena via stimulation of mu, delta, and kappa opioid receptors that regulate

stress perception, pain control, reward behavior, and

neurohormone secretion in reward-relevant brain systems. It has the highest affinity for mu, followed by delta and kappa. Rapid activation of the mu opioid receptor by morphine results in a euphoric phenotype, thus conferring the reinforcing effects of the drug. This activation is accompanied by extracellular dopamine release, which alters several events related to the cAMP signal transduction pathway. Of particular significance is that CREB seems to be modified by morphine, thereby affecting addictive behavioral phenomena, such as withdrawal symptoms.

Applications: ELISA (1:5000-10000)

Reactivity: Species independent

techsupport@bioss.com.cn

Predicted MW.: 0.75883 kDa

400-901-9800