

**bs-2912R****[ Primary Antibody ]****LC3B Rabbit pAb****BioSS**  
**ANTIBODIES**

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

**— DATASHEET —**

<b>Host:</b> Rabbit <b>Clonality:</b> Polyclonal <b>GeneID:</b> 81631 <b>Target:</b> LC3B <b>Immunogen:</b> KLH conjugated synthetic peptide derived from human LC3A/B: 21-121/121. <b>Purification:</b> affinity purified by Protein A <b>Concentration:</b> 1mg/ml <b>Storage:</b> 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. <b>Background:</b> A major contributor to cellular homeostasis is the ability of the cell to strike a balance between the formation and degradation/removal of its cellular components. This process of internal cellular turn-over is called autophagy (self-eating), and is facilitated by a pathway of around 16 interacting proteins in the human. LC3, a ubiquitin-like modifier protein, is the human homolog of yeast Apg8 and is involved in the formation of autophagosomal vacuoles, called autophagosomes. LC3 is expressed as 3 splice variants (LC3A, LC3B and LC3C), which exhibit different tissue distributions and are processed into cytosolic and autophagosomal membrane-bound forms, termed LC3-I and LC3-II, respectively. A disruption to the autophagic process is now associated with the progression of several cancers, neurodegenerative disorders and cardiac pathologies, where LC3 is widely employed as a marker for autophagy.	<b>Isotype:</b> IgG <b>SWISS:</b> Q9GZQ8 <b>Applications:</b> <b>IHC-P</b> (1:100-500) <b>IHC-F</b> (1:100-500) <b>IF</b> (1:100-500) <b>Reactivity:</b> Mouse, Rat, Rabbit (predicted: Human, Pig, Cow, Zebrafish, Chicken, Dog, Horse) <b>Predicted MW.:</b> 13 kDa <b>Subcellular Location:</b> Cell membrane ,Cytoplasm
---	---

**— VALIDATION IMAGES —**

Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by microwave in sodium citrate buffer (pH6.0) ; Block endogenous peroxidase by 3% hydrogen peroxide for 30 minutes; Blocking buffer (3% BSA) at RT for 30min; Antibody incubation with (LC3B) Polyclonal Antibody, Unconjugated (bs-2912R) at 1:400 overnight at 4°C, followed by conjugation to the secondary antibody (labeled with HRP) and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Rabbit brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (LC3B) Polyclonal Antibody, Unconjugated (bs-2912R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

**— SELECTED CITATIONS —**

- **[IF=15.304]** Jinbo Li. et al. Autophagy inhibition recovers deficient ICD-based cancer immunotherapy. BIOMATERIALS. 2022 Aug;287:121651 IF ;Mouse. 35777331
- **[IF=15.1]** Xingbo Wang. et al. Construction of a cascade nanosystem to implement indirect and direct cell modulation

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

for tumor microenvironment immunostimulation. CHEM ENG J. 2024 Apr;485:150141 WB ;Mouse.  
10.1016/j.cej.2024.150141

- **[IF=13.281]** Jinjin Shi. et al. Photoactivated Self-Disassembly of Multifunctional DNA Nanoflower Enables Amplified Autophagy Suppression for Low-Dose Photodynamic Therapy. 2021 Oct 20 IF ;Mouse. 34672076
- **[IF=10.6]** Chen Jiang. et al. Maternal exposure to nanopolystyrene induces neurotoxicity in offspring through P53-mediated ferritinophagy and ferroptosis in the rat hippocampus. J NANOBIOTECHNOL. 2024 Dec;22(1):1-15 IF ;Rat. 39438901
- **[IF=8.5]** Liu-Lu Gao. et al. Acteoside suppresses hepatocellular carcinoma progression via modulation of macrophage migration inhibitory factor and mitogen-activated protein kinase proteins. INT J BIOL MACROMOL. 2025 Jun;;145579 WB ;Human. 40582652