

bs-3281R**[Primary Antibody]****Phospho-MSK1 (Ser360) Rabbit pAb****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

— DATASHEET —


Host: Rabbit	Isotype: IgG	Applications: WB (1:500-2000)
Clonality: Polyclonal		IHC-P (1:100-500)
GeneID: 9252	SWISS: O75582	IHC-F (1:100-500)
Target: Phospho-MSK1 (Ser360)		IF (1:100-500)
Immunogen: KLH conjugated synthesised phosphopeptide derived from human MSK1 around the phosphorylation site of Ser360: TY(p-S)PA.		Reactivity: Mouse, Rat (predicted: Human, Rabbit, Cow, Chicken, Dog)
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		Predicted MW.: 90 kDa
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		Subcellular Location: Cytoplasm ,Nucleus
Background: MSK1 is a mitogen and stress activated protein kinase 1 which belongs to the AGC family of kinases and is related in structure to the ribosomal p70 S6 kinase subfamily. MSK1 can be activated by ERK1/2 and SAPK2/p38 MAP kinase. It is also known to be required for the phosphorylation of CREB, ATF1 H3 and HMG14 in response to mitogen and stress. Similar to RSK, MSK1 contains two kinase domains (N term and a C term). Once phosphorylated on Thr581 and Ser360 by ERK1/2 and SAPK2/p38, MSK1 autophosphorylate on at least 5 sites. Of these autophosphorylation sites Ser212 and Ser376 get phosphorylated by the C terminal kinase domain of MSK1 which is essential for the catalytic activity of the N terminal kinase domain.		

— VALIDATION IMAGES —

Sample: Lymph node lysates(mo); Primary: Rabbit Anti-Phospho-MSK1 (Ser360) (bs-3281R) at 1:300 dilution; Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size:90 kD Observed band size:100 kD



Tissue/cell: Mouse brain tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Incubation: Anti-MSK1 Polyclonal Antibody, Unconjugated(bs-3281R) 1:500, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Paraformaldehyde-fixed, paraffin embedded (Rat kidney); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Phospho-MSK1 (Ser360)) Polyclonal Antibody, Unconjugated (bs-3281R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.