bs-3282R

[Primary Antibody]

phospho-MSK1 (Ser212) Rabbit pAb



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

GenelD: 9252 **SWISS:** 075582

Target: MSK1 (Ser212)

Immunogen: KLH conjugated synthesised phosphopeptide derived from human

MSK1 around the phosphorylation site of Ser212: AY(p-S)FC.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: MSK1 is a mitogen and stress activated protein kinase 1 which

belongs to the AGC family of kinases and is related in structure to the ribosomal p70 S6 kinase subfamily. MSK1 can be activated by ERK1/2 and SAPK2/p38 MAP kinase. It is also known to be required for the phosphorylation of CREB, ATF1 H3 and HMG14 in response to mitogen and stress. Similar to RSK, MSK1 contains two kinase domains (N term and a C term). Once phosphorylated on Thr581 and Ser360 by ERK1/2 and SAPK2/p38, MSK1 autophosphorylate on at least 5 sites. Of these autophosphorylation sites Ser212 and Ser376 get phosphorylated by the C terminal kinase domain of MSK1 which is essential for the catalytic activity of the N terminal kinase domain.

Applications: WB (1:500-2000)

IHC-P (1:100-500) IHC-F (1:100-500) **IF** (1:100-500) Flow-Cyt (2ug/Test) ICC/IF (1:100-500)

Reactivity: Human, Rat


(predicted: Mouse, Rabbit, Pig, Cow, Chicken, Dog,

Horse)


Predicted 90 kDa

Subcellular Location: Cytoplasm ,Nucleus


VALIDATION IMAGES




Sample: Lane 1: Human 293T cell lysates Lane 2: Human HeLa cell lysates Lane 3: Human MCF-7 cell lysates Lane 4: Human HUVEC cell lysates Lane 5: Human A549 cell lysates Lane 6: Human MDA-MB-231 cell lysates Primary: Anti-Phospho-MSK1 (Ser212) (bs-3282R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 90 kDa Observed band size: 85 kDa



Tissue/cell: Rat brain tissue; 4% Paraformaldehyde-fixed and paraffinembedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Incubation: Anti-phospho-MSK1 Polyclonal Antibody. Unconjugated(bs-3282R) 1:500, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Hela cell; 4% Paraformaldehyde-fixed; Triton X-100 at room temperature for 20 min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Antibody incubation with (Phospho-MSK1 (Ser212)) polyclonal Antibody, Unconjugated (bs-3282R) 1:50, 90 minutes at 37°C; followed by a conjugated Goat Anti-Rabbit IgG antibody at 37°C for 90 minutes, DAPI (blue, C02-04002) was used to stain the cell nuclei.



Blank control (black line) :Hela. Primary Antibody (green line): Rabbit Anti-PhosphoMSK1(Ser212) antibody (bs-3282R)
Dilution:2ug/Test; Secondary Antibody (white blue line): Goat anti-rabbit IgG-FITC Dilution: 0.5ug/Test. Isotype control (orange line): Normal Rabbit IgG Protocol The cells were fixed with 4% PFA (10min at room temperature) and then permeabilized with 90% ice-cold methanol for 20 min at -20°C, The cells were then incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at room temperature. Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.