

bs-6583R**[Primary Antibody]****RNASE4 Rabbit pAb****BioSS**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

— DATASHEET —

Host: Rabbit Clonality: Polyclonal GeneID: 6038 Target: RNASE4 Purification: affinity purified by Protein A Concentration: 1mg/ml Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. Background: The protein encoded by this gene belongs to the pancreatic ribonuclease family. It plays an important role in mRNA cleavage and has marked specificity towards the 3' side of uridine nucleotides. Alternative splicing results in four transcript variants encoding the same protein. This gene and the gene that encodes angiogenin share promoters and 5' exons. Each gene splices to a unique downstream exon that contains its complete coding region. [provided by RefSeq, Aug 2013].	Isotype: IgG SWISS: P34096	Applications: WB (1:500-2000) IHC-P (1:100-500) Reactivity: Mouse, Rat (predicted: Human, Rabbit, Dog, Horse) Predicted MW.: 14 kDa Subcellular Location: Secreted
--	---	---

— VALIDATION IMAGES —

Sample: Lane 1: Human HepG2 cell lysates Lane 2: Human HUVEC cell lysates
 Primary: Anti-RNASE4 (bs-6583R) at 1/1000 dilution
 Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution
 Predicted band size: 14 kDa
 Observed band size: 15 kDa



Tissue/cell: rat lung tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Incubation: Anti-RNASE4/Angiogenin Polyclonal Antibody, Unconjugated(bs-6583R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining