- DATASHEET -

Host: Rabbit

Clonality: Polyclonal

GenelD: 2892

[Primary Antibody]

Glutamate receptor 3 Rabbit pAb



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

Isotype: IgG SWISS: P42263 Applications: IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) IF (1:100-500) Reactivity: Human (predicted: Mouse, Rat, Rabbit, Pig, Cow, Dog, Horse) Predicted MW.: 98 kDa

Subcellular Extracellular matrix ,Cell Location: membrane

Immunogen: KLH conjugated synthetic peptide derived from human GLUR3: 451-550/894. < Extracellular >

Purification: affinity purified by Protein A

Target: Glutamate receptor 3

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: L-Glutamate is the major excitatory neurotransmitter in the mammalian CNS, acting through both ligand gated ion channels (ionotropic receptors) and G-protein coupled (metabotropic) receptors. As such the glutamate receptors play a vital role in the mediation of excitatory synaptic transmission. The ionotropic glutamate receptors are multimeric assemblies of four or five subunits, and are subdivided into three groups (AMPA, NMDA and Kainate receptors) based on their pharmacology structural similarities Glutamate Receptor 3 is a ionotropic, AMPA-selective, membrane bound protein widely distributed in the CNS. Predominant expression is seen in the neuronal cells of the cerebral cortex, dentate gyrus, and glial cells throughout brain regions. AMPA receptors mediate fast synaptic transmission in the CNS and are composed of subunits GluR1-4, products from separate genes. Glutamate Receptor 3 exists in 2 named isoforms -Flip and Flop, produced by alternative splicing.

– VALIDATION IMAGES



Paraformaldehyde-fixed, paraffin embedded (Human breast cancer); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Glutamate receptor 3) Polyclonal Antibody, Unconjugated (bs-1799R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.