## bs-5497R

## [ Primary Antibody ]

www.bioss.com.cn

sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

DATASHEET -

Host: Rabbit Isotype: IgG

phospho-MAPK3 (Tyr204) Rabbit pAb

Clonality: Polyclonal

Target: MAPK3 (Tyr204)

**Immunogen:** KLH conjugated Synthesised phosphopeptide derived from rat

MAPK3 around the phosphorylation site of Tyr204: TE(p-Y)VA.

**Purification:** affinity purified by Protein A

Concentration: 1mg/ml

**Storage:** 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

**Background:** The protein encoded by this gene is a member of the MAPkinase

family. MAP kinases, also known as extracellularsignal-regulated kinases (ERKs), act in a signaling cascade that regulates various cellular processes such as proliferation, differentiation, and cell cycle progression in response to avariety of extracellular signals. This kinase is activated byupstream kinases, resulting in its translocation to the nucleus where it phosphorylates nuclear targets. Alternatively splicedtranscript variants encoding different

protein isoforms have beendescribed. [provided by RefSeq, Jul 2008].

Applications: IHC-P (1:100-500)

IHC-F (1:100-500) **IF** (1:100-500)

Reactivity: Mouse (predicted: Human,


Rat, Rabbit, Pig, Cow, Chicken, Dog, GuineaPig,

Horse)

**Predicted** 43 kDa MW.:

Subcellular Location: Nucleus

## VALIDATION IMAGES



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (phospho-MAPK3(Tyr204)) Polyclonal Antibody, Unconjugated (bs-5497R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.