

bs-5503R**[Primary Antibody]****phospho-MAPKAPK2 (Ser272) Rabbit pAb****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

— DATASHEET —


Host: Rabbit	Isotype: IgG	Applications: WB (1:500-2000)
Clonality: Polyclonal		IHC-P (1:100-500)
GeneID: 9261	SWISS: P49137	IHC-F (1:100-500)
Target: MAPKAPK2 (Ser272)		IF (1:100-500)
Immunogen: KLH conjugated Synthesised phosphopeptide derived from human MAPKAPK2 around the phosphorylation site of Ser272: AI(p-S)PG.		Reactivity: Human, Mouse, Rat (predicted: Rabbit, Pig, Cow, Chicken)
Purification: affinity purified by Protein A		Predicted MW.: 46 kDa
Concentration: 1mg/ml		Subcellular Location: Cytoplasm ,Nucleus
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: MAP kinase activated protein kinase 2 (MAPKAP Kinase 2), also known as p45 hsp27 kinase, is a 45-54 kDa serine/threonine protein kinase that contains a proline rich sequence and two putative SH3 binding sites. MAPKAP Kinase 2 is activated in response to stress, IL1 and TNF, possibly catalyzed by p38/Hog dependent phosphorylation. One of the major substrates of MAPKAP Kinase 2 is hsp27, which stimulates actin polymerization in order to facilitate recovery from destruction of cytoskeleton during cellular stresses. MAPKAP2 is implicated in several disorders including ischemic brain injury and heart failure and has been shown to be important in regulating stress resistance and the production of TNF alpha.		

— VALIDATION IMAGES —

Sample: Lung (Mouse) Lysate at 40 ug Primary: Anti-Phospho-MAPKAPK2(Ser272) (bs-5503R) at 1/200 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 46 kD Observed band size: 47 kD



Paraformaldehyde-fixed, paraffin embedded (mouse skeletal muscle); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Incubation with (Phospho-MAPKAPK2 (Ser272)) Polyclonal Antibody, Unconjugated (bs-5503R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (rat skeletal muscle); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Incubation with (Phospho-MAPKAPK2 (Ser272)) Polyclonal Antibody, Unconjugated (bs-5503R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

— SELECTED CITATIONS —

- **[IF=2.72]** Shafieyan, Yousef, et al. "Effects of Low Frequency Cyclic Mechanical Stretching on Osteoclastogenesis." Journal of Biomechanics (2014). WB ;Mouse. /10.1016/j.jbiomech.2014.06.028