

bs-24941R**[Primary Antibody]****MYH7 Rabbit pAb****BioSS**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

— DATASHEET —


Host: Rabbit	Isotype: IgG	Applications: IHC-P (1:400-800) IHC-F (1:400-800) IF (1:100-500)
Clonality: Polyclonal		
GeneID: 140781	SWISS: Q91Z83	
Target: MYH7		Reactivity: Mouse, Rat (predicted: Human, Sheep, Dog)
Immunogen: KLH conjugated synthetic peptide derived from mouse MYH7: 1870-1935/1935.		
Purification: affinity purified by Protein A		Predicted MW.: 213 kDa
Concentration: 1mg/ml		Subcellular Location: Cytoplasm
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: Myosin heavy chains are ubiquitous Actin-based motor proteins that convert the chemical energy derived from ATP hydrolysis into the mechanical energy that drives diverse motile processes in eukaryotic cells, including cytokinesis, vesicular transport and cellular locomotion. Muscle myosin is a heterohexamer consisting of two myosin heavy chains and two associated nonidentical pairs of myosin light chains. The seven myosin heavy chain isoforms that predominate in mammalian skeletal muscles include two developmental isoforms, MHC-embryonic (MYH3) and MHC-perinatal (MYH8); three adult skeletal muscle isoforms, MHC IIa (MYH2), MHC IIb (MYH4) and MHC IIx/d (MYH1); and MHC- β /slow (MYH7 or MHC- β), which is also expressed in cardiac muscle. Research indicates that mutations of the MYH7 gene causes hypertrophic cardiomyopathy.		

— VALIDATION IMAGES —

Paraformaldehyde-fixed, paraffin embedded (mouse skeletal muscle); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (MYH7) Polyclonal Antibody, Unconjugated (bs-24941R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructionsand DAB staining.



Paraformaldehyde-fixed, paraffin embedded (rat skeletal muscle); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (MYH7) Polyclonal Antibody, Unconjugated (bs-24941R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructionsand DAB staining.



Paraformaldehyde-fixed, paraffin embedded (rat skeletal muscle); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (MYH7) Polyclonal Antibody, Unconjugated (bs-20941R) at 1:200 overnight at 4°C, followed by a conjugated Goat Anti-Rabbit IgG antibody (bs-0295G-FITC) for 90 minutes, and DAPI for nuclei staining.

— SELECTED CITATIONS —

- **[IF=2.2]** Wenqi Ding. et al. Changes of mRNA, miRNA and lncRNA expression contributing to skeletal muscle differences between fetus and adult Mongolian horses. COMP BIOCHEM PHYS D. 2024 Jul;101294 IHC ;Horse. 10.1016/j.cbd.2024.101294

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.