

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

MC1R Rabbit pAb

Catalog Number: bs-23516R

Target Protein: MC1R
Concentration: 1mg/ml

Form: Liquid

Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: WB (1:500-2000)

Reactivity: Mouse, Rat

Predicted MW: 35 kDa

Entrez Gene: 17199

Source: KLH conjugated synthetic peptide derived from mouse MC1 Receptor: 231-315/315.

Purification: affinity purified by Protein A

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: Enables melanocyte-stimulating hormone receptor activity. Involved in intracellular signal

transduction; positive regulation of intracellular signal transduction; and positive regulation

of transcription by RNA polymerase II. Acts upstream of or within melanin biosynthetic

process; pigmentation; and sensory perception of pain. Predicted to be located in

membrane. Predicted to be integral component of membrane. Predicted to be active in

cytoplasm and plasma membrane. Is expressed in ductus deferens; epididymis; esophagus;


and skin. Human ortholog(s) of this gene implicated in familial melanoma; major depressive

 $\ disorder; melanoma; oculocutaneous \ albinism \ type \ II; and \ pigmentation \ disease.$


Orthologous to human MC1R (melanocortin 1 receptor). [provided by Alliance of Genome

Resources, Nov 2021]

VALIDATION IMAGES



Sample: B16(Mouse) Cell Lysate at 30 ug Primary: Anti-MC1 Receptor (bs-23516R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 35 kD Observed band size: 45 kD



Sample: Adrenal gland (Rat) Lysate at 40 ug Primary: Anti-MC1 Receptor (bs-23516R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 35 kD Observed band size: 45 kD