bs-3787R

[Primary Antibody]

Bioss

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

SCD Rabbit pAb

- DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

GenelD: 6319 **SWISS:** 000767

Target: SCD

Immunogen: KLH conjugated synthetic peptide derived from human SCD:

265-359/359.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: This gene encodes an enzyme involved in fatty acid

biosynthesis, primarily the synthesis of oleic acid. The protein belongs to the fatty acid desaturase family and

is an integral membrane protein located in the endoplasmic reticulum. Transcripts of approximately

3.9 and 5.2 kb, differing only by alternative polyadenlyation signals, have been detected. A gene encoding a similar enzyme is located on chromosome 4

and a pseudogene of this gene is located on chromosome 17. [provided by RefSeq, Sep 2015]

Applications: IHC-P (1:100-500)

IHC-F (1:100-500) IF (1:100-500) ICC/IF (1:100-500)


Reactivity: Human, Mouse, Rat

(predicted: Rabbit, Pig, Sheep, Cow, Dog)

Predicted MW.: 41 kDa

Subcellular Cell membrane, Cytoplasm


VALIDATION IMAGES



Paraformaldehyde-fixed, paraffin embedded (rat spleen); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (SCD1) Polyclonal Antibody, Unconjugated (bs-3787R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (human colon carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (SCD1) Polyclonal Antibody, Unconjugated (bs-3787R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



A431 cell; 4% Paraformaldehyde-fixed; Triton X-100 at room temperature for 20 min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Antibody incubation with (SCD1) polyclonal Antibody, Unconjugated (bs-3787R) 1:100, 90 minutes at 37°C; followed by a conjugated Goat Anti-Rabbit IgG antibody at 37°C for 90 minutes, DAPI (blue, C02-04002) was used to stain the cell nuclei

- SELECTED CITATIONS -

- [IF=9.8] Yanghuan Yu. et al. MiRNA-seq and mRNA-seq revealed the mechanism of fluoride-induced cauda epididymal injury. SCI TOTAL ENVIRON. 2024 Jun;930:172895 WB; Mouse. 38697545
- [IF=9.685] Wang, Nan. et al. USP7- and PRMT5-dependent G3BP2 stabilization drives de novo lipogenesis and tumorigenesis of HNSC. CELL DEATH DIS. 2023 Mar;14(3):1-13 WB; Human. 36878903

- [IF=7.5] Jing Zhang. et al. Influence of coated cobalamin on lactation performance, nutrient digestion, and proteins related to mammary epithelial cell proliferation and milk fatty acid synthesis in dairy cows. ANIM NUTR. 2025 Aug;: WB ;Bovine. 10.1016/j.aninu.2025.06.006
- [IF=6.1] Jiangtao Huang. et al. Enhancement of PPARα-Inhibited Leucine Metabolism-Stimulated β-Casein Synthesis and Fatty Acid Synthesis in Primary Bovine Mammary Epithelial Cells. J AGR FOOD CHEM. 2023;XXXX(XXX):XXX-XXX WB;Bovine. 37853551
- [IF=6.3] Jing Zhang. et al. Dietary supplementation of sodium butyrate enhances lactation performance by promoting nutrient digestion and mammary gland development in dairy cows. ANIM NUTR. 2023 Sep;: WB; Bovine. 10.1016/j.aninu.2023.08.008