

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

Desmoglein 3 Rabbit pAb

Catalog Number: bs-6585R

Target Protein: Desmoglein 3

Concentration: 1mg/ml

Form: Liquid

Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: IHC-P (1:100-500), IHC-F (1:100-500), IF (1:100-500)

Reactivity: Mouse, Rat (predicted: Human, Rabbit, Dog)

Predicted MW: 105 kDa Entrez Gene: 1830

Swiss Prot: P32926

Source: KLH conjugated synthetic peptide derived from human Desmoglein 3: 50-75/999.

Purification: affinity purified by Protein A

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: Desmosomes are cell-cell junctions between epithelial, myocardial, and certain other cell

types. Desmoglein 3 is a calcium-binding transmembrane glycoprotein component of


desmosomes in vertebrate epithelial cells. Currently, three desmoglein subfamily members

have been identified and all are members of the cadherin cell adhesion molecule superfamily. These desmoglein gene family members are located in a cluster on

chromosome 18. This protein has been identified as the autoantigen of the autoimmune

skin blistering disease pemphigus vulgaris.

VALIDATION IMAGES



Tissue/cell: rat tongue tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min; Incubation: Anti-Desmoglein 3 Polyclonal Antibody, Unconjugated(bs-6585R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining

PRODUCT SPECIFIC PUBLICATIONS

[IF=4.996] Oike, Asami. et al. Bisphosphonate affects the behavioral responses to HCl by disrupting farnesyl diphosphate synthase in mouse taste bud and tongue epithelial cells. SCI REP-UK. 2022 Dec;12(1):1-15 IF; MOUSE . 36481783

[IF=4.3] Sumitomo et al. Streptococcal Cysteine Protease-Mediated Cleavage of Desmogleins Is Involved in the Pathogenesis of Cutaneous Infection. (2018) Front.Cell.Infect.Microbiol. 8:10 IF; MOUSE . 29416987

[IF=0.32] BİR, Ferda, et al. "New immunohistochemical markers in the differential diagnosis of nonsmall cell lung carcinoma." Turkish Journal of Medical Sciences (2016). IHC; = "Human". doi:10.3906/sag-1501-68