

---

## Beta casein Rabbit pAb

Catalog Number: bs-23551R

Target Protein: Beta casein

Concentration: 1mg/ml

Form: Liquid

Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: WB (1:500-2000)

Reactivity: Human, Cow, Goat

Predicted MW: 24 kDa

Entrez Gene: 1447

Swiss Prot: P05814

Source: KLH conjugated synthetic peptide derived from human Beta casein: 1-100/226.

Purification: affinity purified by Protein A


Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

**Background:** Milk proteins are crucial for the development of all newborn mammals and caseins constitute the major proteins in mammalian milk. b- and k-caseins are the only caseins present in human milk. The b-casein/k-casein ratio is higher in colostrum than in transitional and mature milk and is related to a better digestibility of colostrum casein micelles by the neonate during the first days of life. Human b-casein-encoding gene (Bca) contains a highly phosphorylated site, which is responsible for the calcium-binding capacity of b-casein. A common set of transcription factors are required for the expression of b-casein. Multiple binding sites for Stat5, C/EBPb (CCAAT/enhancer-binding protein) and several half-sites for glucocorticoid receptor (GR) are identified in the distal human enhancer of the b-casein gene. b-casein gene transcription is regulated primarily by a composite response element (CoRE), which integrates signaling from the lactogenic hormones PRL, insulin and hydrocortisone in mammary epithelial cells. NFkB functions as a negative regulator of b-casein gene expression during pregnancy by interfering with Stat5 tyrosine phosphorylation

### VALIDATION IMAGES

---



Sample: Lane 1: Milk (Goat) Lysate at 2 ug Lane 2: Milk (Cow) Lysate at 30 ug Lane 3: Milk (Cow) Lysate at 3 ug Lane 4: Milk (Human) Lysate at 30 ug Lane 5: Milk (Human) Lysate at 3 ug Primary: Anti-Beta-casein (bs-23551R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 25-29 kD Observed band size: 25-29 kD

## PRODUCT SPECIFIC PUBLICATIONS

**[IF=3.7]** Wei Yu. et al. Glucose promotes cell growth and casein synthesis via ATF4/Nrf2-Sestrin2- AMPK-mTORC1 pathway in dairy cow mammary epithelial cells. ANIM BIOTECHNOL. 2023 Jul 12 WB ; Bovine . 37435839