
BRN3B Rabbit pAb

Catalog Number: bs-6985R

Target Protein: BRN3B

Concentration: 1mg/ml

Form: Liquid

Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: IHC-P (1:100-500), IHC-F (1:100-500), IF (1:50-200), Flow-Cyt (1µg /Test)

Reactivity: Mouse, Rat (predicted:Human, Pig, Sheep, Cow, Dog, Horse)

Predicted MW: 41 kDa

Entrez Gene: 5458

Swiss Prot: Q12837.2

Source: KLH conjugated synthetic peptide derived from human BRN3B: 301-400/409.


Purification: affinity purified by Protein A

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.


Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: The Brn family of transcription factors are found in a highly restricted subset of neurons and are critical to the early embryonic development of the central nervous system. Brn-1 and Brn-2 are class III POU (Pit-Oct-Unc) domain proteins, whereas Brn-3 is a class IV POU domain protein. Three Brn-3 proteins have been described and are designated Brn-3a, Brn-3b and Brn-3c. While Brn-3a and Brn-3c stimulate transcription, Brn-3b generally functions as a transcriptional repressor. However, Brn-3b, but not Brn-3a, has been shown to regulate the expression of the acetylcholine receptor. Interestingly, Brn-3a has two functional transactivating domains, one at the amino-terminus and one at the carboxy-terminus. Brn-2 is thought to be involved in smooth muscle cell development and differentiation.

VALIDATION IMAGES



Paraformaldehyde-fixed, paraffin embedded (mouse testis); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (BRN3B) Polyclonal Antibody, Unconjugated (bs-6985R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Blank control: RSC96(fixed with 2% paraformaldehyde (10 min) and then permeabilized with ice-cold 90% methanol for 30 min on ice). Primary Antibody: Rabbit Anti-BRN3B antibody(bs-6985R), Dilution: 1μg in 100 μL 1X PBS containing 0.5% BSA; Isotype Control Antibody: Rabbit IgG(orange) ,used under the same conditions); Secondary Antibody: Goat anti-rabbit IgG-PE(white blue), Dilution: 1:200 in 1 X PBS containing 0.5% BSA.

PRODUCT SPECIFIC PUBLICATIONS

[IF=2.1] Zeng Qi. et al. TRIM9 promotes Müller cell-derived retinal stem cells to differentiate into retinal ganglion cells by regulating Atoh7. IN VITRO CELL DEV-AN. 2023 Oct;;1-10 IF ; Rat . 37792226