

bs-5335R**[Primary Antibody]****Phospho-ELk1 (Ser389) Rabbit pAb****BioSS**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

— DATASHEET —

Host: Rabbit	Isotype: IgG	Applications: WB (1:500-2000) Flow-Cyt (1ug/Test)
Clonality: Polyclonal		
GeneID: 2002	SWISS: P19419	Reactivity: Human, Mouse, Rat (predicted: Rabbit, Pig, Sheep, Cow, Dog, GuineaPig)
Target: Phospho-ELk1 (Ser389)		Predicted MW.: 47 kDa
Immunogen: KLH conjugated Synthesised phosphopeptide derived from human ELK1 around the phosphorylation site of Ser389: PR(p-S)PA.		Subcellular Location: Nucleus
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: This gene is a member of the Ets family of transcription factors and of the ternary complex factor (TCF) subfamily. Proteins of the TCF subfamily form a ternary complex by binding to the the serum response factor and the serum response element in the promoter of the c-fos proto-oncogene. The protein encoded by this gene is a nuclear target for the ras-raf-MAPK signaling cascade. This gene produces multiple isoforms by using alternative translational start codons and by alternative splicing. Related pseudogenes have been identified on chromosomes 7 and 14. [provided by RefSeq, Mar 2012].		

— VALIDATION IMAGES —

Sample: Lane 1: Mouse Cerebrum tissue lysates
Lane 2: Rat Cerebrum tissue lysates Primary:
Anti-Phospho-ELk1 (Ser389) (bs-5335R) at
1/1000 dilution Secondary: IRDye800CW Goat
Anti-Rabbit IgG at 1/20000 dilution Predicted
band size: 47 kDa Observed band size: 59 kDa



Blank control (black line) :U-87MG. Primary
Antibody (green line): Rabbit Anti-Phospho-ELk1
(Ser389) antibody (bs-5335R) Dilution:1ug/Test;
Secondary Antibody (white/blue line) : Goat
anti-rabbit IgG-AF488 Dilution: 0.5ug/Test.
Isotype control (orange line) : Normal Rabbit
IgG Protocol The cells were fixed with 4% PFA
(10min at room temperature)and then
permeabilized with 90% ice-cold methanol for
20 min at -20°C, The cells were then incubated in
5%BSA to block non-specific protein-protein
interactions for 30 min at room temperature
.Cells stained with Primary Antibody for 30 min
at room temperature. The secondary antibody
used for 40 min at room temperature.
Acquisition of 20,000 events was performed.

— SELECTED CITATIONS —

- **[IF=7.9]** Shuang Yang, et al. Kanglexin counters vascular smooth muscle cell dedifferentiation and associated arteriosclerosis through inhibiting PDGFR. PHYTOMEDICINE. 2024 May;;155704 WB ;Mouse,Rat. 38759316

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.