bs-55139R

[Primary Antibody]

BIOSS ANTIBODIES

www.bioss.com.cn sales@bioss.com.cn

techsupport@bioss.com.cn 400-901-9800

– DATASHEET –

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

NCOA2 Rabbit pAb

GenelD: 10499 **SWISS:** Q15596

Target: NCOA2

Immunogen: Recombinant human NCOA2: 550-820/1424.

Concentration: 1mg/ml

Storage: PBS with 0.02% Proclin300,50% Glycerol,pH7.3.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: Steroid and thyroid hormones and retinoic acid regulate a complex

array of gene expression activity via intracellular receptor transcription factors belonging to the ligand dependent nuclear receptor superfamily. Adding to the complexity of function of these transcription factors are associated proteins known as coactivators and corepressors which, as their names suggest, enhance or depress transcriptional activity of the nuclear receptor with which they associate. One such coactivator is KAT13C / nuclear receptor coactivator 2 (NCOA2), also termed Glucocorticoid receptor-

interacting protein 1 (GRIP1).

Applications: WB (1:500-2000)

IHC-P (1:100-500) IHC-F (1:400-800) IF (1:100-500) ELISA (1:5000-10000)

Reactivity: Human, Rat

Predicted MW.: 159 kDa

Subcellular Nucleus