
phospho-Tau (Thr205) Rabbit pAb

Catalog Number: bs-5420R

Target Protein: phospho-Tau (Thr205)

Concentration: 1mg/ml

Form: Liquid

Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: WB (1:500-2000), IHC-P (1:100-500), IHC-F (1:100-500), IF (1:100-500)

Reactivity: Human, Mouse, Rat (predicted:Rabbit, Dog, Horse)

Predicted MW: 52/79 kDa

Entrez Gene: 4137

Swiss Prot: P10636

Source: KLH conjugated Synthesised phosphopeptide derived from human Tau around the phosphorylation site of Thr205: SR(p-T)PS.


Purification: affinity purified by Protein A

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.


Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: Tau proteins are important Promotes microtubule assembly and stability, and might be involved in the establishment and maintenance of neuronal polarity. The C-terminus binds axonal microtubules while the N-terminus binds neural plasma membrane components, suggesting that tau functions as a linker protein between both. Axonal polarity is predetermined by tau localization (in the neuronal cell) in the domain of the cell body defined by the centrosome. The short isoforms allow plasticity of the cytoskeleton whereas the longer isoforms may preferentially play a role in its stabilization. Tau proteins subcellular located in the axons of neurons, in the cytosol and in association with plasma membrane components. It expressed in neurons. PNS-tau is expressed in the peripheral nervous system while the others are expressed in the central nervous system.


VALIDATION IMAGES



Sample: BV-2(Mouse) Cell Lysate at 30 ug Primary: Anti-phospho-Tau (Thr205) (bs-5420R) at 1/1000 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 52/79 kD Observed
band size: 52/129 kD



Sample: Cerebrum (Mouse) Lysate at 40 ug Cerebrum (Rat) Lysate at 40 ug Primary: Anti-phospho-Tau
(Thr205) (bs-5420R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution
Predicted band size: 52/79 kD Observed band size: 52 kD



Sample: Cerebrum (Mouse) Lysate at 40 ug Cerebrum (Rat) Lysate at 40 ug Cerebellum (Mouse) Lysate at 40
ug Cerebellum (Rat) Lysate at 40 ug Primary: Anti-phospho-Tau (Th205) (bs-5420R) at 1/1000 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 50-70 kD Observed
band size: 50 kD

PRODUCT SPECIFIC PUBLICATIONS

[IF=2.1] Yong Fan. et al. The effect of AQP4 on tau protein aggregation in neurodegeneration and persistent neuroinflammation after cerebral microinfarcts. OPEN MED-WARSAW. 2023 Jan;18(1): WB,IF ; Mouse . 37873537