

bs-1449R**[Primary Antibody]**

Bioss
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

Prohibitin Rabbit pAb**— DATASHEET —**


Host: Rabbit	Isotype: IgG	Applications: WB (1:500-2000) IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) Flow-Cyt (2ug/test) ICC/IF (1:100)
Clonality: Polyclonal		
GeneID: 5245	SWISS: P35232	
Target: Prohibitin		
Immunogen: KLH conjugated synthetic peptide derived from human Prohibitin: 171-272/272.		
Purification: affinity purified by Protein A		Reactivity: Human, Mouse, Rat (predicted: Dog, Horse)
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: This gene is evolutionarily conserved, and its product is proposed to play a role in human cellular senescence and tumor suppression. Antiproliferative activity is reported to be localized to the 3' UTR, which is proposed to function as a trans-acting regulatory RNA. Several pseudogenes of this gene have been identified. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jul 2013]		Predicted MW.: 30 kDa Subcellular Location: Cytoplasm

— VALIDATION IMAGES —


Sample: Heart(Mouse) Lysate at 40 ug Primary: Anti-Prohibitin (bs-1449R) at 1/300 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 30 kD
Observed band size: 30 kD




Sample: A431(Human) Cell Lysate at 40 ug Primary: Anti-Prohibitin (bs-1449R) at 1/300 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 30 kD
Observed band size: 30 kD



Tissue/cell: Human kidney tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Incubation: Anti-Prohibitin Polyclonal Antibody, Unconjugated(bs-1449R) 1:500, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Tissue/cell: Rat intestine tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Incubation:



HepG2 cell; 4% Paraformaldehyde-fixed; Triton X-100 at room temperature for 20 min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Antibody incubation with (Prohibitin) polyclonal Antibody, Unconjugated (bs-1449R) 1:100, 90 minutes at 37°C; followed by a conjugated Goat Anti-Rabbit IgG antibody at

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Anti-Prohibitin Polyclonal Antibody,
Unconjugated(bs-1449R) 1:500, overnight at 4°C,
followed by conjugation to the secondary
antibody(SP-0023) and DAB(C-0010) staining

37°C for 90 minutes, DAPI (blue, C02-04002) was
used to stain the cell nuclei.

— SELECTED CITATIONS —

- **[IF=5.23]** Wang, Jinzi, et al. "Comparative Secretome Analysis Reveals Perturbation of Host Secretion Pathways by a Hypovirus." Scientific Reports 6 (2016): 34308. WB ;="Other Species". 27698384
- **[IF=2.08]** Xing, Wen Min, et al. "Proteomic identification of mitochondrial targets involved in andrographolide sodium bisulfite-induced nephrotoxicity in a rat model." Environmental Toxicology and Pharmacology (2015). WB ;="Rat". 26356389