

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

phospho-heterochromatin protein 1 (pTyr24) Rabbit pAb

Catalog Number: bs-0928R

Target Protein: phospho-heterochromatin protein 1 (pTyr24)

Concentration: 1mg/ml

Form: Liquid
Host: Rabbit
Clonality: Polyclonal

Isotype: IgG

Applications: WB (1:500-2000), IHC-P (1:100-500), IHC-F (1:100-500), IF (1:100-500), ELISA (1:5000-10000)

Reactivity: (predicted:Fruit Fly)

Predicted MW: 23 kDa

Source: KLH conjugated Synthesised phosphopeptide derived from Fruit Fly HP1 around the

phosphorylation site of Tyr24: EE(p-Y)AV.

Purification: affinity purified by Protein A

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: HP1 alpha is a component of heterochromatin. It recognizes and binds histone H3 tails

methylated at Lys-9, leading to epigenetic repression. HP1 alpha may interact with lamin B receptor (LBR). This interaction can contribute to the association of the heterochromatin with the inner nuclear membrane. HP1 proteins are relatively small proteins (~25 kDa) with a conserved amino-terminal chromo domain and a structurally related carboxy-terminal motif, the chromo shadow domain. Both domains of HP1 are required for binding to native chromatin in vivo, but they contribute differentially to binding in euchromatin and

chromatin in vivo, but they contribute unierentially to binding in euclifornatin and

heterochromatin.

PRODUCT SPECIFIC PUBLICATIONS

[IF=3.057] Terzuoli et al. Antagonism of bradykinin B2 receptor prevents inflammatory responses in human endothelial cells by quenching the NF-kB pathway activation. (2014) PLoS.On. 9:e84358 WB; Human . 24392129