## bs-5424R

## [ Primary Antibody ]

## BIOSS ANTIBODIES

www.bioss.com.cn

sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**GenelD:** 257397 **SWISS:** Q8N5C8

Target: MAP3K7IP3

**Immunogen:** KLH conjugated synthetic peptide derived from human

MAP3K7IP3: 601-712/712.

**Purification:** affinity purified by Protein A

MAP3K7IP3 Rabbit pAb

Concentration: 1mg/ml

**Storage:** 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

**Background:** The product of this gene functions in the NF-kappaB signal

transduction pathway. The encoded protein, and the similar and functionally redundant protein MAP3K7IP2/TAB2, forms a ternary complex with the protein kinase MAP3K7/TAK1 and either TRAF2 or TRAF6 in response to stimulation with the pro-inflammatory cytokines TNF or IL-1. Subsequent MAP3K7/TAK1 kinase activity triggers a signaling cascade leading to activation of the NF-kappaB transcription factor. The human genome contains a related pseudogene. Alternatively spliced transcript variants have been described, but their biological validity has not been determined.

Applications: WB (1:500-2000)

IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500)

**ELISA** (1:5000-10000)

Reactivity: Rat (predicted: Human,

Mouse, Rabbit, Pig, Sheep,

Cow, Dog, Horse)

Predicted MW.: 79 kDa

**Subcellular** Extracellular matrix ,Cell **Location:** membrane ,Cytoplasm

## - SELECTED CITATIONS -

• [IF=6.43] Hussain et al. MicroRNA 27a-3p Regulates Antimicrobial Responses of Murine Macrophages Infected byMycobacterium aviumsubspeciesparatuberculosisby Targeting Interleukin-10 and TGF-β-Activated Protein Kinase 1 Binding Protein 2. (2018) Front.Immunol. 8:1915 WB; Mouse. 29375563