

bs-8453R**[Primary Antibody]****phospho-Tau (Thr181) Rabbit pAb****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

— DATASHEET —

Host: Rabbit Clonality: Polyclonal GeneID: 4137 Target: Tau (Thr181) Immunogen: KLH conjugated Synthesised phosphopeptide derived from human Tau around the phosphorylation site of Thr181: PK(p-T)PP. Purification: affinity purified by Protein A Concentration: 1mg/ml Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. Background: Tau proteins are important Promotes microtubule assembly and stability, and might be involved in the establishment and maintenance of neuronal polarity. The C-terminus binds axonal microtubules while the N-terminus binds neural plasma membrane components, suggesting that tau functions as a linker protein between both. Axonal polarity is predetermined by tau localization (in the neuronal cell) in the domain of the cell body defined by the centrosome. The short isoforms allow plasticity of the cytoskeleton whereas the longer isoforms may preferentially play a role in its stabilization. Tau proteins subcellular located in the axons of neurons, in the cytosol and in association with plasma membrane components. It expressed in neurons. PNS-tau is expressed in the peripheral nervous system while the others are expressed in the central nervous system.	Isotype: IgG SWISS: P10636 Applications: WB (1:500-2000) Reactivity: Human, Mouse, Rat (predicted: Rabbit, Cow, Dog, Horse) Predicted MW.: 52/83 kDa Subcellular Location: Cell membrane ,Cytoplasm
---	---

— VALIDATION IMAGES —

25 ug total protein per lane of various lysates
(see on figure) probed with phospho-Tau
(Thr181) polyclonal antibody, unconjugated
(bs-8453R) at 1:1000 dilution and 4°C overnight
incubation. Followed by conjugated secondary
antibody incubation at r.t. for 60 min.