bs-8460R

[Primary Antibody]

FNDC1 Rabbit pAb

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal GeneID: 84624 Target: FNDC1

Immunogen: KLH conjugated synthetic peptide derived from human FNDC1:

21-120/1894.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: FNDC1, also known as MEL4B3, Ags8, Expressed in synovial lining protein and Activation-associated cDNA protein, is a 1888 amino

acid secreted protein that contains five fibronectin type-III domains. FNDC1 is moderately expressed in skeletal muscle, pancreas, heart, kidney, spinal cord, ovary and lung. Expression of

FNDC1 is induced in response to hypoxia in ventricular

cardiomyocytes. Since FNDC1 interacts with G ∫ and G©, it is likely that FNDC1 is an activator for G-protein signaling. Though normally absent in healthy skin, FNDC1 expression is induced by TGF-beta signaling in skin tumors and psoriasis. There are two isoforms of FNDC1 that are produced as a result of alternative splicing events.

Applications: IHC-P (1:100-500)

IHC-F (1:100-500) **IF** (1:50-200)

ELISA (1:5000-10000)

Reactivity: (predicted: Human, Mouse,

Rat, Rabbit, Sheep, Cow,

Dog, Horse)

Predicted MW.: 202 kDa

Subcellular Location: Secreted

— SELECTED CITATIONS —

- [IF=9.4] ZhangRui Xin. et al. FNDC1 is a myokine that promotes myogenesis and muscle regeneration. EMBO J. 2024 Nov 20 WB, IF, CoIP; Mouse. 39567831
- [IF=4.848] Jiang T et al. FNDC1 Promotes the Invasiveness of Gastric Cancer via Wnt/β-Catenin Signaling Pathway and Correlates With Peritoneal Metastasis and PrognosisFront Oncol.2020 Dec 17;10:590492. IHC; Human. 33392086