## bs-1450R

- DATASHEET -

## [ Primary Antibody ]

Isotype: IgG

SWISS: P03986

## TRGC2 Rabbit pAb

Host: Rabbit

Clonality: Polyclonal

GenelD: 6965



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

Applications: WB (1:500-2000) **IHC-P** (1:100-500) **IHC-F** (1:100-500) IF (1:100-500)

Reactivity: Mouse, Rat (predicted: Human, Cow, Horse)

Predicted MW.: 22/35 kDa

Subcellular Location: Cell membrane

Target: TRGC2 **Immunogen:** KLH conjugated synthetic peptide derived from human TCRG: 1-100/189. Purification: affinity purified by Protein A Concentration: 1mg/ml Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. Background: T cell receptors recognize foreign antigens which have been processed as small peptides and bound to major histocompatibility complex (MHC) molecules at the surface of

antigen presenting cells (APC). Each T cell receptor is a dimer consisting of one alpha and one beta chain or one delta and one gamma chain. In a single cell, the T cell receptor loci are rearranged and expressed in the order delta, gamma, beta, and alpha. If both delta and gamma rearrangements produce functional chains, the cell expresses delta and gamma. If not, the cell proceeds to rearrange the beta and alpha loci. This region represents the germline organization of the T cell receptor gamma locus. The gamma locus includes V (variable), J (joining), and C (constant) segments. During T cell development, the gamma chain is synthesized by a recombination event at the DNA level joining a V segment with a J segment; the C segment is later joined by splicing at the RNA level. Recombination of many different V segments with several J segments provides a wide range of antigen recognition. Additional diversity is attained by junctional diversity, resulting from the random addition of nucleotides by terminal deoxynucleotidyltransferase. Several V segments of the gamma locus are known to be incapable of encoding a protein and are considered pseudogenes. Somatic rearrangement of the gamma locus has been observed in T cells derived from patients with T cell leukemia and ataxia telangiectasia. [provided by RefSeq, Jul 2008]

## - VALIDATION IMAGES



Sample: Bone (mouse) Lysate at 40 ug Primary: Anti- TCRG(bs-1450R)at 1/300 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 22/35kD Observed band size: 27/40 kD



Tissue/cell: rat spleen tissue; 4% Paraformaldehyde-fixed and paraffinembedded: Antigen retrieval: citrate buffer ( 0.01M, pH 6.0 ), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min; Incubation: Anti-TCRG Polyclonal Antibody. Unconjugated(bs-1450R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining