

bs-9183R**[Primary Antibody]****VGLL2 Rabbit pAb****BioSS**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

— DATASHEET —

Host: Rabbit	Isotype: IgG	Applications: WB (1:500-2000)
Clonality: Polyclonal		Reactivity: Mouse (predicted: Human, Rat, Rabbit, Pig, Cow, Dog)
GeneID: 245806	SWISS: Q8N8G2	
Target: VGLL2		Predicted MW.: 33 kDa
Immunogen: KLH conjugated synthetic peptide derived from human VGLL2: 101-200/317.		Subcellular Location: Nucleus
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: Vgl-2, also known as VITO-1, is a 317 amino acid protein that contains a domain through which it interacts with TEF-1, a protein that plays a role in controlling the expression of numerous genes. Specific to skeletal muscle, Vgl-2 is expressed highly in adult fast muscle and is expressed at lower levels in adult slow muscle and fetal skeletal muscle. During muscle differentiation, Vgl-2 mRNA levels increase and Vgl-2 translocates from the cytoplasm to the nucleus. Overexpression of Vgl-2 in MYOD-transfected 10T1/2 mouse embryonic fibroblasts increases expression of myosin heavy chain (MHC), which is a marker of terminal muscle differentiation. This evidence suggests that Vgl-2 is essential for muscle gene expression. There are two isoforms of Vgl-2 that are produced as a result of alternative splicing events.		

— VALIDATION IMAGES —

Sample: Muscle (Mouse) Lysate at 40 ug Primary:
Anti-VGLL2 (bs-9183R) at 1/300 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at
1/20000 dilution Predicted band size: 33 kD
Observed band size: 35 kD