

**bsm-52646R****[ Primary Antibody ]****TNFRSF12A/CD266 Recombinant Rabbit mAb****Bioss**  
**ANTIBODIES**

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

**— DATASHEET —**

<b>Host:</b> Rabbit	<b>Isotype:</b> IgG	<b>Applications:</b> WB (1:500-2000)  <b>Reactivity:</b> Human, Mouse (predicted: Rat)  <b>Predicted MW.:</b> 11 kDa  <b>Subcellular Location:</b> Cell membrane ,Cytoplasm
<b>Clonality:</b> Recombinant	<b>CloneNo.:</b> 2C14	
<b>GeneID:</b> 51330	<b>SWISS:</b> Q9NP84	
<b>Target:</b> TNFRSF12A/CD266		
<b>Immunogen:</b> A synthesized peptide derived from human CD266: 81-129.		
<b>Purification:</b> affinity purified by Protein A		
<b>Concentration:</b> 1mg/ml		
<b>Storage:</b> 1*TBS (pH7.4), 0.05% BSA, 40% Glycerol. Preservative: 0.02% Proclin300. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
<b>Background:</b> TWEAK (otherwise known as TNFRSF12A, Tumor necrosis factor receptor superfamily member 12A), is a member of the TNF family and was first described as a weak inducer of apoptosis in some cell types. It is thought to promote angiogenesis and the proliferation of endothelial cells as well as modulating cellular adhesion to matrix proteins. Recently, a receptor for TWEAK was isolated by expression cloning from a HUVEC cell cDNA library. Function : Receptor for TNFSF12/TWEAK. Weak inducer of apoptosis in some cell types. Promotes angiogenesis and the proliferation of endothelial cells. May modulate cellular adhesion to matrix proteins.		

**— VALIDATION IMAGES —**

25 ug total protein per lane of various lysates (see on figure) probed with TNFRSF12A/CD266 monoclonal antibody, unconjugated (bsm-52646R) at 1:1000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min.