bs-24530R

[Primary Antibody]

phospho-SMAD2 (Ser464) Rabbit pAb



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

GenelD: 4087 **SWISS:** Q15796

Target: SMAD2 (Ser464)

Immunogen: KLH conjugated synthesised phosphopeptide derived from human

Smad2 around the phosphorylation site of Ser464: RC(p-S)SM.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: The protein encoded by this gene belongs to the SMAD, a family of proteins similar to the gene products of the Drosophila gene 'mothers against decapentaplegic' (Mad) and the C. elegans gene Sma. SMAD proteins are signal transducers and transcriptional modulators that mediate multiple signaling pathways. This protein mediates the signal of the transforming growth factor (TGF)-beta, and thus regulates multiple cellular processes, such as cell proliferation, apoptosis, and differentiation. This protein is recruited to the TGF-beta receptors through its interaction with the SMAD anchor for receptor activation (SARA) protein. In response to TGF-beta signal, this protein is phosphorylated by the TGF-beta receptors. The phosphorylation induces the dissociation of this protein with SARA and the association with the family member SMAD4. The association with SMAD4 is important for the translocation of this protein into the nucleus, where it binds to target promoters and forms a transcription repressor complex with other cofactors. This protein can also be phosphorylated by activin type 1 receptor kinase, and mediates the signal from the activin. Alternatively spliced transcript variants have been observed for this gene. [provided by RefSeq, May 2012]

Applications: WB (1:500-2000)

IHC-P (1:100-500) **IHC-F** (1:100-500) **IF** (1:100-500) Flow-Cyt (3ug/Test)

Reactivity: Human, Mouse


(predicted: Rat, Cow, Chicken, Dog, Horse)

Predicted 58 kDa


MW.:

Location: Cytoplasm ,Nucleus Subcellular


VALIDATION IMAGES




Sample: Lane1: Heart (Mouse) Lysate at 40 ug Lane2: Muscle (Mouse) Lysate at 40 ug Primary: Anti-phospho-SMAD2 (Ser464) (bs-24530R) at 1/300 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 58 kD Observed band size: 58 kD




Sample: Lane 1: Hela (Human) Cell Lysate at 30 ug Lane 2: HT1080 (Human) Cell Lysate at 30 ug Lane 3: Jurkat (Human) Cell Lysate at 30 ug Lane 4: A549 (Human) Cell Lysate at 30 ug Lane 5: 293T (Human) Cell Lysate at 30 ug Primary: Antiphospho-SMAD2 (Ser464) (bs-24530R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 60 kD Observed band size: 60 kD



Paraformaldehyde-fixed, paraffin embedded (human placenta); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (phospho-SMAD2 (Ser464)) Polyclonal Antibody, Unconjugated (bs-24530R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (phospho-SMAD2 (Ser464)) Polyclonal Antibody, Unconjugated (bs-24530R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Blank control (Black line): Mouse spleen(Black). Primary Antibody (green line): Rabbit Antiphospho-SMAD2(Ser425) antibody (bs-0457R) Dilution: 3µg/10^6 cells; Isotype Control Antibody (orange line): Rabbit IgG . Secondary Antibody (white blue line): Goat anti-rabbit IgG-PE Dilution: 1µg /test. Protocol The cells were fixed with 4% PFA (10min at room temperature) and then permeabilized with 90% ice-cold methanol for 20 min at room temperature. The cells were then incubated in 5%BSA goat serum to block non-specific protein-protein interactions for 15 min at room temperature .Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.

- SELECTED CITATIONS -

• [IF=6.656] Leihao Hu. et al. Identification of the Active Compounds in the Yi-Fei-San-Jie Formula Using A Comprehensive Strategy Based on Cell Extraction/UPLC-MS/MS, Network Pharmacology, and Molecular Biology Techniques. PHYTOMEDICINE. 2023 Apr;:154843 WB; Human. 37149966