
Phospho-CHEK2 (Ser516) Rabbit pAb

Catalog Number: bs-5257R

Target Protein: Phospho-CHEK2 (Ser516)

Concentration: 1mg/ml

Form: Liquid

Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: WB (1:500-2000)

Reactivity: Human (predicted: Mouse, Rat, Rabbit, Pig, Chicken, Dog)

Predicted MW: 61 kDa

Subcellular: Nucleus

Locations:

Entrez Gene: 11200

Swiss Prot: O96017

Source: KLH conjugated Synthesised phosphopeptide derived from human CHEK2 around the phosphorylation site of Ser516: QP(p-S)TS.


Purification: affinity purified by Protein A

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.


Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: In response to DNA damage and replication blocks, cell cycle progression is halted through the control of critical cell cycle regulators. The protein encoded by this gene is a cell cycle checkpoint regulator and putative tumor suppressor. It contains a forkhead-associated protein interaction domain essential for activation in response to DNA damage and is rapidly phosphorylated in response to replication blocks and DNA damage. When activated, the encoded protein is known to inhibit CDC25C phosphatase, preventing entry into mitosis, and has been shown to stabilize the tumor suppressor protein p53, leading to cell cycle arrest in G1. In addition, this protein interacts with and phosphorylates BRCA1, allowing BRCA1 to restore survival after DNA damage. Mutations in this gene have been linked with Li-Fraumeni syndrome, a highly penetrant familial cancer phenotype usually associated with inherited mutations in TP53. Also, mutations in this gene are thought to confer a predisposition to sarcomas, breast cancer, and brain tumors. This nuclear protein is a member of the CDS1 subfamily of serine/threonine protein kinases. Several transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Apr


VALIDATION IMAGES



Sample: HeLa UV(Human) Cell Lysate at 30 ug HeLa (Human) Cell Lysate at 30 ug Primary: Anti-p-Chk2(Ser516)(bs-5257R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 61 kD Observed band size: 56 kD



Sample: 293T(Human) Cell Lysate at 30 ug Primary: Anti-Phospho-Chk2 (Ser516) (bs-5257R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 61 kD Observed band size: 57 kD



Sample: HeLa(Human) Cell Lysate at 30 ug Primary: Anti-Phospho-Chk2 (Ser516) (bs-5257R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 61 kD Observed band size: 57 kD