

bs-24389R**[Primary Antibody]****FUT8 Rabbit pAb****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

— DATASHEET —

Host: Rabbit	Isotype: IgG	Applications: WB (1:500-2000)
Clonality: Polyclonal		Reactivity: Mouse, Rat (predicted: Human, Rabbit, Chicken, Dog, Horse)
GeneID: 2530	SWISS: Q9BYC5	Predicted MW.: 66 kDa
Target: FUT8		Subcellular Location: Cell membrane ,Cytoplasm
Immunogen: KLH conjugated synthetic peptide derived from human FUT8 : 201-300/575.		
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: Catalyzes the addition of fucose in alpha 1-6 linkage to the first GlcNAc residue, next to the peptide chains in N-glycans. The expression of this gene may contribute to the malignancy of cancer cells and to their invasive and metastatic capabilities. There are 2 isoforms produced by alternative splicing. Immunogen is not present in isoform 2 (also known as Retinal).		

— VALIDATION IMAGES —

Sample: Lane 1: Lung (Mouse) Lysate at 40 ug
Lane 2: Lung (Rat) Lysate at 40 ug Primary: Anti-FUT8 (bs-24389R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 66 kD Observed band size: 70 kD

— SELECTED CITATIONS —

- **[IF=7.4]** Wen Zhang. et al. FUT8-Mediated Core Fucosylation Promotes the Pulmonary Vascular Remodeling in Pulmonary Arterial Hypertension. AGING DIS. 2023 Oct 1; 14(5): 1927-1944 IF,WB ;Rat. 37196106