- DATASHEET -

[Primary Antibody]

Histone H4 (Acetyl K8) Rabbit pAb



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

| Host: Rabbit | Isotype: IgG | |
|---------------------------------|-------------------------------------|---|
| Clonality: Polyclonal | | |
| GenelD: 121504 | SWISS: P62805 | |
| Target: Histone H4 (Acetyl K8) | | |
| munogon: KI H conjugated Synthe | sised acetuloeptide derived from hu | ~ |

Immunogen: KLH conjugated Synthesised acetylpeptide derived from human Histone H4 around the acetylation site of Lys8: GG(Ac-K)GL.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a member of the histone H4 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. [provided by RefSeq, Jul 2008]

– VALIDATION IMAGES



Sample: Lane 1: Human Hela cell lysates Lane 2: Human Jurkat cell lysates Lane 3: Human HL60 cell lysates Lane 4: Human MCF-7 cell lysates Primary: Anti-Histone H4 (Acetyl K8) (bs-4016R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 11 kD Observed band size: 11 kD



Paraformaldehyde-fixed, paraffin embedded (Rat liver); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Histone H4 (Acetyl K8)) Polyclonal Antibody, Unconjugated (bs-4016R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Rat skin); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Histone H4 (Acetyl K8)) Polyclonal Antibody, Unconjugated (bs-4016R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Hela cell; 4% Paraformaldehyde-fixed; Triton X-100 at room temperature for 20 min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20



Blank control (black line) :NIH/3T3. Primary Antibody (green line): Rabbit Anti-Histone H4 (Acetyl K8) antibody (bs-4016R)

Applications: WB (1:500-2000) IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) Flow-Cyt (1ug/test)

ICC/IF (1:100)

Reactivity: Human, Mouse, Rat (predicted: Rabbit, Pig, Cow, Dog, Horse)

Predicted MW.: 11 kDa

Subcellular Location: Nucleus min; Antibody incubation with (Histone H4 (Acetyl K8)) polyclonal Antibody, Unconjugated (bs-4016R) 1:100, 90 minutes at 37°C; followed by a conjugated Goat Anti-Rabbit IgG antibody at 37°C for 90 minutes, DAPI (blue, C02-04002) was used to stain the cell nuclei. Dilution:1ug/Test; Secondary Antibody (white blue line) : Goat anti-rabbit IgG-AF488 Dilution: 0.5ug/Test. Isotype control (orange line) : Normal Rabbit IgG Protocol The cells were fixed with 4% PFA (10min at room temperature)and then permeabilized with 90% ice-cold methanol for 20 min at -20°C, The cells were then incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at room temperature .Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.