

bs-3901R**[Primary Antibody]****Cubilin Rabbit pAb****BioSS**
ANTIBODIES

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

— DATASHEET —

Host: Rabbit	Isotype: IgG	Applications: IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) ELISA (1:5000-10000) Reactivity: Mouse (predicted: Human, Rat) Predicted MW.: 395 kDa Subcellular Location: Cell membrane ,Cytoplasm
Clonality: Polyclonal		
GeneID: 8029	SWISS: O60494	
Target: Cubilin		
Immunogen: KLH conjugated synthetic peptide derived from human Cubilin: 51-150/3623.		
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: Cubilin is an endocytic receptor that lacks a classical transmembrane region. It is a multidomain receptor that contains an amino terminal 110 residue segment followed by 8 epidermal growth factor (EGF)-like repeats and a contiguous stretch of 27 CUB domains. Cubilin acts as a receptor for intrinsic factor-vitamin B12 complexes. Cubilin is located within the epithelium of intestine and kidney. Cubilin co-localizes with and binds to Megalin, a 600kDa member of the LDL receptor family that is required for the internalization of cubilin-bound ligands, such as vitamin B12, apolipoprotein A1 and Albumin. Megalin specifically binds to cubilin in the amino terminal region that contains the EGF-like repeats and CUB domains 1 and 2. Mutations in Cubilin may play a role in autosomal recessive megaloblastic anemia also known as Imlerslund-Grasbeck's disease, which causes intestinal malabsorption of vitamin B12.		