bs-3943R

[Primary Antibody]


GNB3 Rabbit pAb



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

| - DATASHEET | |
|---|--|
| Isotype: IgG | Applications: IHC-P (1:100-500) IHC-F (1:100-500) |
| SWISS: P16520 | IF (1:100-500) |
| | Reactivity: Human (predicted: Mouse, Rat, Rabbit, Pig, Sheep, Cow, Dog, Horse) |
| synthetic peptide derived from human G protein 0. | |
| y Protein A | Predicted MW.: ^{37 kDa} |
| | |
| l) with 1% BSA, 0.02% Proclin300 and 50% tore at -20°C for one year. Avoid repeated es. | Subcellular Cell membrane ,Cytoplasm Location: |
| alpha subunits and for G protein-effector ngle-nucleotide polymorphism (C825T) in this d with essential hypertension and obesity and is | |
| | SWISS: P16520 Synthetic peptide derived from human G protein 0. y Protein A 4) with 1% BSA, 0.02% Proclin300 and 50% tore at -20°C for one year. Avoid repeated es. mposed of 3 units, alpha, beta and gamma and e modulation and transduction of various signalling systems. There are families of related bunit. GNB3 encodes a beta subunit. Beta irred for GTPase activity, for replacement of GDP alpha subunits and for G protein-effector ngle-nucleotide polymorphism (C825T) in this d with essential hypertension and obesity and is <i>i</i> th the occurrence of the splice variant GNB3-s, |

- VALIDATION IMAGES -



Paraformaldehyde-fixed, paraffin embedded (human cerebellum); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Incubation with (FXYD1) Polyclonal Antibody, Unconjugated (bs-3943R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructionsand DAB staining.