bs-0979R

[Primary Antibody]

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

CD38 Rabbit pAb

DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Target: CD38

Immunogen: KLH conjugated synthetic peptide derived from mouse CD38:

101-200/304. < Extracellular >

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: The protein encoded by this gene is a non-lineage-restricted, type II transmembrane glycoprotein that synthesizes and hydrolyzes cyclic adenosine 5'-diphosphate-ribose, an intracellular calcium ion mobilizing messenger. The release of soluble protein and the ability of membrane-bound protein to become internalized indicate both extracellular and intracellular functions for the protein. This protein has an N-terminal cytoplasmic tail, a single membrane-spanning domain, and a C-terminal extracellular region with four N-glycosylation sites. Crystal structure analysis demonstrates that the functional molecule is a dimer, with the central portion containing the catalytic site. It is used as a prognostic marker for patients with chronic lymphocytic leukemia. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Sep 2015]

Applications: WB (1:500-2000)

IHC-P (1:100-500) **IHC-F** (1:100-500) **IF** (1:100-500)

Reactivity: Mouse, Rat


(predicted: Human)

Predicted


34 kDa MW.:

Subcellular Cell membrane

VALIDATION IMAGES



Sample: Cerebrum (Rat) Lysate at 40 ug Primary: Anti-CD38 (bs-0979R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 34 kD Observed band size: 48 kD



Sample: Pancreas (Mouse) Lysate at 40 ug Primary: Anti-CD38 (bs-0979R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 34 kD Observed band size: 48 kD



Tissue/cell: rat brain tissue; 4% Paraformaldehyde-fixed and paraffinembedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Incubation: Anti-CD38 Polyclonal Antibody, Unconjugated(bs-0979R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining

- SELECTED CITATIONS -

- [IF=14.919] Jinfeng Wang. et al. Translocation of vaginal microbiota is involved in impairment and protection of uterine health. Nat Commun. 2021 Jul;12(1):1-15 IF; Rat. 34234149
- [IF=8.9] Guo Yang. et al. Mechanisms of electroacupuncture-induced neuroprotection in acute stroke rats: the role of astrocyte-mediated mitochondrial transfer. CELL COMMUN SIGNAL. 2025 Dec;23(1):1-20 WB ;Rat. 40598228

[IF=4.966] Santoshi Muppalaet al. The P387 thrombospondin - 4 variant promotes accumulation of macrophages in atherosclerotic lesions. FASEB J. 2020 Sep;34(9):11529-11545. IHC,IF; mouse. 32686880
[IF=1.257] Moridera K et al. Skeletal unloading reduces cluster of differentiation (CD) 38 expression in the bone marrow and osteoblasts of mice. J Orthop Sci. 2019 May 6. ICC; Mouse. 31072650