

bs-5432R**[Primary Antibody]****phospho-MEK5 (Ser137) Rabbit pAb****BioSS**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

— DATASHEET —

| | | |
|---|----------------------|--|
| Host: Rabbit | Isotype: IgG | Applications: WB (1:500-2000) |
| Clonality: Polyclonal | | Reactivity: Human, Mouse (predicted: Rabbit, Sheep, Cow, Horse) |
| GeneID: 5607 | SWISS: Q13163 | |
| Target: MEK5 (Ser137) | | Predicted MW.: 49 kDa |
| Immunogen: KLH conjugated Synthesised phosphopeptide derived from human MEK5 around the phosphorylation site of Ser137: AV(p-S)DS. | | Subcellular Location: Cytoplasm |
| Purification: affinity purified by Protein A | | |
| Concentration: 1mg/ml | | |
| Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. | | |
| Background: MEK5 is a dual specificity protein kinase belonging to the Ser/Thr protein kinase family, (MAP kinase kinase family). It is activated by phosphorylation on Ser/Thr by MAP kinase kinases and interacts specifically with ERK5, and not with another MAP kinase like P38. This kinase specifically interacts with and activates MAPK7/ERK5. This kinase itself can be phosphorylated and activated by MAP3K3/MEKK3, as well as by atypical protein kinase C isoforms (aPKCs). It is not phosphorylated by RAFA, RAFB or RAFC and it may interact with GTPases such as CDC42. The signal cascade mediated by this kinase is involved in growth factor stimulated cell proliferation and muscle cell differentiation. MEK5 is expressed in many adult tissues and is most abundant in heart and skeletal muscle. | | |

— VALIDATION IMAGES —

Sample: MDA-MB-231 Cell (Human) Lysate at 40 ug
Primary: Anti-phospho-MEK5(Ser137)
(bs-5432R) at 1/300 dilution Secondary:
IRDye800CW Goat Anti-Rabbit IgG at 1/20000
dilution Predicted band size: 49 kD Observed
band size: 50 kD



Sample: Cerebrum (Mouse) Lysate at 40 ug
Primary: Anti-phospho-MEK5(Ser137)
(bs-5432R) at 1/300 dilution Secondary:
IRDye800CW Goat Anti-Rabbit IgG at 1/20000
dilution Predicted band size: 49 kD Observed
band size: 50 kD