bs-13801R

[Primary Antibody]

C1GALT1C1 Rabbit pAb



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

| | 400-901-9800 |
|---|---|
| Isotype: IgG | Applications: WB (1:500-2000) |
| | IHC-P (1:100-500) IHC-F (1:100-500) |
| SWISS: Q96EU7 | IF (1:100-500) |
| | ICC/IF (1:100-500) |
| 0/318. Protein A with 1% BSA, 0.02% Proclin300 and 50% re at -20°C for one year. Avoid repeated a type II transmembrane protein that is 3-galactosyltransferase 1, which catalyz e-1 structure, also known as Thomsen- n, on O-linked glycans. This gene produ ferase activity itself, but instead acts as ne required for the folding, stability and 1 beta1,3-galactosyltransferase 1. Mutar n associated with Tn syndrome. Alternal ariants encoding the same protein have | MW.: ^{36 KDa} Subcellular Location: Cell membrane uct lacks a I full tions in tively |
| | SWISS: Q96EU7 hthetic peptide derived from human 0/318. Protein A with 1% BSA, 0.02% Proclin300 and 50% re at -20°C for one year. Avoid repeated a type II transmembrane protein that is 3-galactosyltransferase 1, which cataly re-1 structure, also known as Thomsen- in, on O-linked glycans. This gene produ ferase activity itself, but instead acts as ne required for the folding, stability and 1 beta1,3-galactosyltransferase 1. Muta n associated with Tn syndrome. Alterna |