

bs-5600R**[Primary Antibody]****phospho-TSC1 (Ser505) Rabbit pAb****BioSS**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

— DATASHEET —

Host: Rabbit Clonality: Polyclonal GeneID: 7248 Target: TSC1 (Ser505) Immunogen: KLH conjugated Synthesised phosphopeptide derived from human TSC1 around the phosphorylation site of Ser505: FD(p-S)PF. Purification: affinity purified by Protein A Concentration: 1mg/ml Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. Background: Tuberous sclerosis complex (TSC) is an autosomal dominant genetic disorder characterized by mental retardation and the widespread development of distinctive tumors termed hamartomas. Two different genetic loci have been linked to TSC; one of these loci, the tuberous sclerosis-2 gene (TSC2), encodes a protein called tuberin and the other loci, tuberous sclerosis-1 gene (TSC1), encodes a protein called hamartin. Tuberin and hamartin interact with each other forming a cytoplasmic complex. Hamartin interacts with the ezrin-radixin-moesin (ERM) family of actin-binding proteins and inhibition of hamartin activity results in loss of cell adhesion. Hamartin is present in most adult tissues with strong expression in brain, heart, and kidney.	Isotype: IgG SWISS: Q92574 Applications: WB (1:500-2000) IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) Reactivity: Human, Mouse (predicted: Rat, Rabbit, Pig, Cow, Chicken, Dog, Horse) Predicted MW.: 128 kDa Subcellular Location: Cell membrane ,Cytoplasm
---	---

— VALIDATION IMAGES —

Sample: NIH/3T3(Mouse) Cell Lysate at 30 ug
Primary: Anti-phospho-TSC1(Ser505) (bs-5600R)
at 1/1000 dilution Secondary: IRDye800CW Goat
Anti-Rabbit IgG at 1/20000 dilution Predicted
band size: 128 kD Observed band size: 125 kD



Paraformaldehyde-fixed, paraffin embedded
(human brain glioma); Antigen retrieval by
boiling in sodium citrate buffer (pH6.0) for
15min; Block endogenous peroxidase by 3%
hydrogen peroxide for 20 minutes; Blocking
buffer (normal goat serum) at 37°C for 30min;
Antibody incubation with (TSC1(Ser505))
Polyclonal Antibody, Unconjugated (bs-5600R)
at 1:400 overnight at 4°C, followed by operating
according to SP Kit(Rabbit) (sp-0023)
instructions and DAB staining.

— SELECTED CITATIONS —

- **[IF=4.8]** Cao, Jiaxue, et al. "DNA methylation Landscape of body size variation in sheep." Scientific reports 5 (2015). WB ;Sheep. 26472088