# bsm-52396R

# [ Primary Antibody ]

# alpha smooth muscle Actin Recombinant Rabbit A N T B =**mAb**

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

### DATASHEET —

Host: Rabbit Isotype: IgG Clonality: Recombinant CloneNo.: 2B10 GeneID: 59 **SWISS:** P62736

Target: alpha smooth muscle Actin **Purification:** affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: All eukaryotic cells express Actin, which often constitutes as much

as 50% of total cellular protein. Actin filaments can form both stable and labile structures and are crucial components of microvilli and the contractile apparatus of muscle cells. While lower eukaryotes, such as yeast, have only one Actin gene, higher eukaryotes have several isoforms encoded by a family of genes. At least six types of Actin are present in mammalian tissues and fall into three classes. alpha-Actin expression is limited to various types of muscle, whereas beta- and gamma-Actin are the principle constituents of filaments in other tissues. Members of the small GTPase family regulate the organization of the Actin cytoskeleton. Rho controls the assembly of Actin stress fibers and focal adhesion. Rac regulates Actin filament accumulation at the plasma

membrane. Cdc42 stimulates formation of filopodia.

Applications: WB (1:1000-5000)

IHC-P (1:100-500) **IHC-F** (1:100-500) **IF** (1:100-500) ICC/IF (1:100-500)


Reactivity: Human, Mouse, Rat

(predicted: Zebrafish)


Predicted MW.: 42 kDa

Subcellular Location: Cytoplasm


# VALIDATION IMAGES




25 ug total protein per lane of various lysates (see on figure) probed with alpha smooth muscle Actin monoclonal antibody. unconjugated (bsm-52396R) at 1:2000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min.




Paraformaldehyde-fixed, paraffin embedded Human Breast Cancer; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with alpha smooth muscle Actin Monoclonal Antibody, Unconjugated(bsm-52396R) at 1:200 overnight at 4°C, followed by conjugation to the bs-0295G-HRP and DAB (C-0010) staining.



Paraformaldehyde-fixed, paraffin embedded Human Uterus; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with alpha smooth muscle Actin Monoclonal Antibody, Unconjugated(bsm-52396R) at 1:200 overnight at 4°C, followed by conjugation to the bs-0295G-HRP and DAB (C-0010) staining.



Paraformaldehyde-fixed, paraffin embedded Human small intestine; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min: Antibody incubation with alpha smooth muscle Actin Monoclonal Antibody, Unconjugated (bsm-52396R) at 1:200 overnight



Paraformaldehyde-fixed, paraffin embedded Mouse stomach; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with alpha smooth muscle Actin Monoclonal Antibody, Unconjugated (bsm-52396R) at 1:200 overnight at 4°C.

at 4°C. Followed by conjugated Goat Anti-Rabbit IgG antibody (green, bs-0295G-BF488), DAPI
(blue, C02-04002) was used to stain the cell

C02-04002) was used to stain the cell nuclei. (blue, C02-04002) was used to stain the cell

Followed by conjugated Goat Anti-Rabbit IgG C02-04002) was used to stain the cell nuclei.

## - SELECTED CITATIONS -

- [IF=17.1] Lu Tan. et al. Mechanically Robust Hemostatic Hydrogel Membranes with Programmable Strain-Adaptive Microdomain Entanglement for Wound Treatment in Dynamic Tissues. ACS NANO. 2024;XXXX(XXX):XXX-XXX IHC;Rat. 38457334
- [IF=11.8] Huanliang Liu. et al. PM2.5 triggers autophagic degradation of Caveolin-1 via endoplasmic reticulum stress (ERS) to enhance the TGF-β1/Smad3 axis promoting pulmonary fibrosis. ENVIRON INT. 2023 Nov;181:108290 WB; Rat. 10.1016/j.envint.2023.108290
- [IF=11.205] Yingchun Luo. et al. Akkermansia muciniphila prevents cold-related atrial fibrillation in rats by modulation of TMAO induced cardiac pyroptosis. EBIOMEDICINE. 2022 Aug;82:104087 WB;Rat. 35797768
- [IF=7.675] Ya-Ling Yin. et al. Citronellal Attenuates Oxidative Stress–Induced Mitochondrial Damage through TRPM2/NHE1 Pathway and Effectively Inhibits Endothelial Dysfunction in Type 2 Diabetes Mellitus. ANTIOXIDANTS-BASEL. 2022 Nov;11(11):2241 IF ;Rat. 36421426
- [IF=5.923] Chih-Hsin Hsu. et al. miR-29a-3p/THBS2 Axis Regulates PAH-Induced Cardiac Fibrosis. Int J Mol Sci. 2021 Jan;22(19):10574 WB,IHC; Human. 10.3390/ijms221910574