bs-23306R

[Primary Antibody]

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn

Semaphorin 3F Rabbit pAb

DATASHEET

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

GeneID: 6405 **SWISS:** Q13275

Target: Semaphorin 3F

Immunogen: KLH conjugated synthetic peptide derived from human Sema3F:

201-300/785.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: The semaphorins are a family of proteins that are involved in signaling. All the family members have a secretion signal, a 500amino acid sema domain, and 16 conserved cysteine residues(Kolodkin et al., 1993 [PubMed 8269517]). Sequence comparisons have grouped the secreted semaphorins into 3 general classes, all of which also have an immunoglobulin domain. The semaphorin III family, consisting of human semaphorin III (SEMA3A; MIM 603961), chicken collapsin, and mouse semaphorins A, D, and E, all have a basic domain at the C terminus. Chicken collapsin contributes to path finding by axons during development by inhibiting extension of growth cones (Luo et al., 1993 [PubMed 8402908]) through an interaction with a collapsin response mediator protein of relative molecular mass 62K (CRMP62) (Goshima et al., 1995 [PubMed7637782]), a putative homolog of an axonal guidance associated UNC33 gene product (MIM 601168). SEMA3F is a secreted member of the semaphorin III family.

Applications: WB (1:500-2000)

400-901-9800


Reactivity: Mouse (predicted: Human,

Rat, Rabbit, Pig, Sheep, Cow, Chicken, Dog)

Predicted MW.: 88 kDa

Subcellular Secreted Location:

VALIDATION IMAGES -



Sample: Lung (Mouse) Lysate at 40 ug Cerebrum (Mouse) Lysate at 40 ug Primary: Anti-Semaphorin 3F (bs-23306R) at 1/1000 dilution Secondary: IRDve800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 88 kD Observed band size: 88 kD

- SELECTED CITATIONS -

• [IF=7.5] Hee Ra Park. et al. Antiseizure effects of Lilii Bulbus on pentylenetetrazol kindling-induced seizures in mice: Involvement of Reelin, Netrin-1, and semaphorin. BIOMED PHARMACOTHER. 2024 Apr;173:116385 WB; Mouse. 38460369