


**bsm-52417R****[ Primary Antibody ]****BioSS**  
**ANTIBODIES**

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

**EpCAM Recombinant Rabbit mAb****— DATASHEET —****Host:** Rabbit**Clonality:** Recombinant**GeneID:** 4072**Target:** EpCAM**Immunogen:** A synthesized peptide derived from human EpCAM: 266-314.**Purification:** affinity purified by Protein A**Concentration:** 1mg/ml**Storage:** 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.  
Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.**Background:** This gene encodes a carcinoma-associated antigen and is a member of a family that includes at least two type I membrane proteins. This antigen is expressed on most normal epithelial cells and gastrointestinal carcinomas and functions as a homotypic calcium-independent cell adhesion molecule. The antigen is being used as a target for immunotherapy treatment of human carcinomas. Mutations in this gene result in congenital tufting enteropathy. [provided by RefSeq, Dec 2008]**Isotype:** IgG**CloneNo.:** 6H5**SWISS:** P16422**Applications:** WB (1:500-2000)**IHC-P** (1:100-500)**IHC-F** (1:100-500)**IF** (1:100-500)**Reactivity:** Human, Mouse, Rat**Predicted MW.:** 35 kDa**Subcellular Location:** Cell membrane**— VALIDATION IMAGES —**


25 ug total protein per lane of various lysates (see on figure) probed with EpCAM monoclonal antibody, unconjugated (bsm-52417R) at 1:2000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min.




Paraformaldehyde-fixed, paraffin embedded (human breast carcinoma); Antigen retrieval by boiling in EDTA buffer (pH9.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Incubation with (EpCAM) Monoclonal Antibody, Unconjugated (bsm-52417R) at 1:100 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (human colon carcinoma); Antigen retrieval by boiling in EDTA buffer (pH9.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Incubation with (EpCAM) Monoclonal Antibody, Unconjugated (bsm-52417R) at 1:100 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded Human Colon; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with EpCAM Monoclonal Antibody, Unconjugated (bsm-52417R) at 1:200 overnight at 4°C. Followed by conjugated Goat Anti-Rabbit IgG antibody (green, bs-0295G-BF488), DAPI (blue, C02-04002) was used to stain



Paraformaldehyde-fixed, paraffin embedded Mouse Colon; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with EpCAM Monoclonal Antibody, Unconjugated (bsm-52417R) at 1:200 overnight at 4°C. Followed by conjugated Goat Anti-Rabbit IgG antibody (green, bs-0295G-BF488), DAPI (blue, C02-04002) was used to stain

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

the cell nuclei.

the cell nuclei.

## — SELECTED CITATIONS —

---

- **[IF=5]** Jun Li. et al. Targeting LRRC41 as a potential therapeutic approach for hepatocellular carcinoma. FRONT MOL BIOSCI. 2023; 10: 1300294 IHC ;Human. 38192337