## bs-5728R

# [ Primary Antibody ]

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

# **HURP Rabbit pAb**

DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**GeneID:** 9787 **SWISS:** Q15398

Target: HURP

**Immunogen:** KLH conjugated synthetic peptide derived from human HURP:

51-150/846.

**Purification:** affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

**Background:** Potential cell cycle regulator that may play a role in carcinogenesis

of cancer cells. Mitotic phosphoprotein regulated by the ubiquitinproteasome pathway. Key regulator of adherens junction integrity and differentiation that may be involved in CDH1-mediated

adhesion and signaling in epithelial cells.

Applications: IHC-P (1:100-500)

IHC-F (1:100-500) **IF** (1:100-500)

Reactivity: Human, Mouse


(predicted: Rat, Rabbit, Pig,

Cow, Dog)

**Predicted** 93 kDa MW.:

Subcellular Cytoplasm ,Nucleus

### VALIDATION IMAGES



Paraformaldehyde-fixed, paraffin embedded (human tonsil); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (HURP) Polyclonal Antibody, Unconjugated (bs-5728R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructionsand DAB staining.



Paraformaldehyde-fixed, paraffin embedded (mouse testis); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (HURP) Polyclonal Antibody, Unconjugated (bs-5728R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.