

bs-3155R**[Primary Antibody]****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

Phospho-GCN2 (Thr899) Rabbit pAb**DATASHEET**

Host: Rabbit Clonality: Polyclonal GeneID: 440275 Target: Phospho-GCN2 (Thr899) Immunogen: KLH conjugated Synthesised phosphopeptide derived from human GCN2 around the phosphorylation site of Thr899: HL(p-T)GM. Purification: affinity purified by Protein A Concentration: 1mg/ml Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. Background: GCN2 belongs to a family of kinases that phosphorylate the alpha subunit of eukaryotic translation initiation factor 2 to downregulate protein synthesis in response to varied cellular stresses.	Isotype: IgG SWISS: Q9P2K8	Applications: Flow-Cyt (1ug/Test) Reactivity: Human, Mouse (predicted: Rat, Cow, Dog, Horse) Predicted MW.: 189 kDa Subcellular Location: Cytoplasm
--	---	---

VALIDATION IMAGES

Blank control (Black line): Molt4 (Black). Primary Antibody (green line): Rabbit Anti-Phospho-GCN2 (Thr899) antibody (bs-3155R) Dilution: 1µg / 10⁶ cells; Isotype Control Antibody (orange line): Rabbit IgG. Secondary Antibody (white/blue line): Goat anti-rabbit IgG-AF647 Dilution: 1µg /test. Protocol The cells were fixed with 4% PFA (10min at room temperature) and then permeabilized with 90% ice-cold methanol for 20 min at room temperature. The cells were then incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at room temperature. Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.



Blank control: Mouse spleen. Primary Antibody (green line): Rabbit Anti-Phospho-GCN2 (Thr899) antibody (bs-3155R) Dilution: 2µg / 10⁶ cells; Isotype Control Antibody (orange line): Rabbit IgG. Secondary Antibody: Goat anti-rabbit IgG-FITC Dilution: 1µg /test. Protocol The cells were fixed with 4% PFA (10min at room temperature) and then permeabilized with 0.1% PBST for 20 min at room temperature. The cells were then incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at room temperature. Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.

SELECTED CITATIONS

- **[IF=25.269]** Changzheng Li. et al. Amino acid catabolism regulates hematopoietic stem cell proteostasis via a GCN2-eIF2α axis. CELL STEM CELL. 2022 Jul;29:1119 WB ;Mouse. 35803229
- **[IF=5.58]** Lehman, Stacey L., Sandra Ryeom, and Constantinos Koumenis. "Signaling through alternative Integrated Stress Response pathways compensates for GCN2 loss in a mouse model of soft tissue sarcoma." Scientific Reports 5

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

(2015). WB ;="Mouse". 26123823

- **[IF=4.9]** Zikang Xing. et al. IDO1 Inhibitor RY103 Suppresses Trp-GCN2-Mediated Angiogenesis and Counters Immunosuppression in Glioblastoma. PHARMACEUTICS. 2024 Jul;16(7):870 WB ;Mouse,Human. 10.3390/pharmaceutics16070870